

PCT/L 00 / 00 33 0
20 AUG 2000

IL00/330

REC'D 29 AUG 2000

WIND

PCT

PA 282142

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

August 07, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 09/329,234

FILING DATE: June 10, 1999

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

W. Montgomery
W. MONTGOMERY
Certifying Officer

JC135 U.S. PTO
09/329234
06/10/99

Sir:

Enclosed herewith for filing is the patent application of

Dr. Mark Friedman
of Shani
Shosegov

**PROCESS OF EXPRESSING AND ISOLATING RECOMBINANT PROTEINS AND RECOMBINANT PROTEIN PRODUCTS FROM
PLANT DERIVED TISSUES OR CULTURED PLANT CELLS**

Enclosed are:

- ☒ 1 sheets of informal drawing(s)
- ☒ An assignment of the invention to CBD TECHNOLOGIES LTD. and to YISSUM RESEARCH AND DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM
- ☐ A certified copy of a _____ application
- ☐ An associate power of attorney
- ☒ A verified statement to establish small entity status under 37 CFR 1.9 and 37 CFR 1.27
- ☐ Other - SEQUENCE LISTING

The filing fee has been calculated as shown below

			SMALL ENTITY		OR	OTHER THAN A SMALL ENTITY	
FOR:	(Col 1) NO FILED	(Col 2) NO. EXTRA	RATE	FEE		RATE	FEE
BASIC FEE				\$ 380	OR		\$
TOTAL CLAIMS	25 - 20 =	5	x9 =	\$ 45	OR	x18 =	\$
INDEP CLAIMS	6 - 3 =	3	x39 =	\$ 117	OR	x78 =	\$
Recordal of Assignment		40	x2 =	\$ 80	OR		\$
* If the difference in Col 1 is less than zero, enter "0" in Col 2			TOTAL	\$ 542	OR	TOTAL	\$

- ☒ Please charge my Deposit Account No. 06-2140 in the amount of \$ 542. A duplicate copy of this sheet is enclosed.
- ☐ A check in the amount of \$ _____ to cover the filing fee is enclosed
- ☒ The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No. 06-2140. A duplicate copy of this sheet is enclosed.
- ☒ Any additional filing fees required under 37 CFR 1.16
- ☒ Any patent application processing fees under 37 CFR 1.17
- ☒ The Commissioner is hereby authorized to charge payment of the following fees during the pendency of this application or credit any overpayment to Deposit Account No. 06-2140. A duplicate copy of this sheet is enclosed.
- ☒ Any patent application processing fees under 37 CFR 1.17.
- ☐ The issue fee set in 37 CFR 1.18 at or before mailing of the Notice of allowance, pursuant to 37 CFR 1.311(b)
- ☒ Any filing fees under 37 CFR 1.16 for presentation of extra claims.

Respectfully,

Mark M. Friedman
Reg. No. 33.883

Mark M. Friedman
DR. MARK FRIEDMAN LTD
C/O Anthony Castorina
2001 Jefferson Davis Highway
Suite 207
Arlington, Virginia 22202

APPLICATION FOR PATENT

5

Inventors: Ziv Shani and Oded Shoseyov

10

Title: PROCESS OF EXPRESSING AND ISOLATING
RECOMBINANT PROTEINS AND RECOMBINANT
PROTEIN PRODUCTS FROM PLANTS, PLANT
DERIVED TISSUES OR CULTURED PLANT CELLS

15

FIELD AND BACKGROUND OF THE INVENTION

20 The present invention relates to a process of expressing and isolating
recombinant proteins and recombinant protein products from plants, plant
derived tissue or cultured plant cells, which process exploits (i) the high
affinity between cellulose binding peptides and cellulose; (ii) the inherent
abundance of cellulose in *planta*; and/or (iii) the simplicity associated with
cellulose isolation from plants, plant derived tissue and cultured plant cells.

25 More particularly, the present invention relates to a process
expressing and isolating recombinant proteins and recombinant protein
products from plants, plant derived tissue or cultured plant cells, which
process employs the expression of a fusion protein including a recombinant
protein and a cellulose binding peptide fused thereto, plant homogenization,
30 isolation of a fusion protein cellulosic matter complex and optional
subsequent isolation of the fusion protein and/or the recombinant protein
from the complex. The present invention further relates to nucleic acid
molecules and to genetically modified or viral infected plants or plant cells
which are useful while implementing the process, and further to a novel
35 composition of matter which results from the process.

Citation or identification of any reference in this section or in any other section of this application shall not be construed as an admission that such reference is available as prior art to the present invention.

5 With the advent of recombinant technology, techniques for the genetic transformation of various host organisms, such as bacteria, yeasts, fungi, plants and animals, for the purposes of producing specific proteins through the expression of heterologous or foreign genes have been extensively developed.

10 Using these recombinant techniques and hosts, numerous commercially important recombinant proteins (examples of which are included hereinbelow) have been expressed and purified. Expression and isolation of a protein of interest on a commercial scale, necessitate the selection of a suitable expression host. This selection largely depends on the economics of production and purification, as well as the ability of the
15 host to accomplish the post-translational modifications needed for full biological activity of the recombinant protein.

Much of the early work in biotechnology was directed toward the expression of recombinant or "heterologous" proteins in prokaryotes like *Escherichia coli* and *Bacillus subtilis* because of the relative ease of genetic
20 manipulation, growth in batch culture and large-scale fermentation of prokaryotes.

Although *E. coli* can in certain cases perform some post translational modifications and events, such as, protein folding and disulfide bond formation, it cannot secrete proteins extracellularly nor can it glycosylate,
25 gamma carboxylate, beta hydroxylate, acetylate or process pre- and pro-peptides. *B. subtilis* suffers from the same limitations as *E. coli* except that it is capable of extracellular secretion.

Furthermore, *E. coli* and other bacteria are pathogens and therefore, depending on the application, contaminants such as pyrogens and

endotoxins expressed along with the recombinant protein must be removed. In addition, extensive post-purification chemical and enzymatic treatments (e.g., to refold the protein into an active form) are sometimes required in order to obtain a biologically active protein.

5 Because proteins are not secreted from prokaryotes like *E. coli*, bacterial cells must be disrupted for product recovery. The subsequent release of bacterial contaminants and other proteins make product purification more difficult and expensive. Because purification accounts for up to 90 % of the total cost of producing recombinant proteins in bacteria, 10 proteins like Insulin can cost several thousand dollars per gram when recombinantly produced in, and subsequently isolated from, *E. coli*.

 Because of the many limitations associated with prokaryotic hosts, the biotechnology industry has looked for eukaryotic host cultures such as, yeast, fungi, insect cells, and mammalian cell tissue culture, to properly and 15 efficiently express recombinant proteins.

 For most of the proteins requiring extensive post-translational modifications for therapeutic and/or functional activity, mammalian cell culture is the most common alternative to *E. coli*. Although mammalian cells are capable of correctly folding and glycosylating bioactive proteins, 20 the quality and extent of glycosylation can vary with different culture conditions among the same host cells. Furthermore, mammalian culture has extremely high fermentation costs (60-80% of total production expense), requires expensive media, and poses safety concerns from potential contamination by viruses and other pathogens. Yields are generally low and 25 in the range of 0.5-1.5% of cellular protein, or micrograms per liter (up to 300-400 milligrams per liter).

 Yeast, other fungi, and insect cells are currently being used as alternatives to mammalian cell culture. Yeast, however, produces incorrectly glycosylated proteins that have excessive mannose residues and

are generally limited in eukaryotic processing. Further, although the baculovirus insect cell system can produce high levels of glycosylated proteins, these are typically not secreted, making purification complex and expensive. Fungi represent the best current system for high-volume, low-cost production of recombinant proteins, but they are not capable of expressing many target proteins.

In addition, eukaryotic cultures, require the maintenance of suitable conditions for efficient commercially viable expression of proteins. As such, the ambient temperature, pH value and aeration level of such cultures need to be carefully controlled, while nutrients must be added to the culture medium in carefully regulated doses and waste products removed. In addition, rigorous aseptic practices must be observed in order to avoid contamination by extraneous microbes. Such cultures are thus normally grown in sophisticated fermentors or bioreactors which are housed in expensively maintained factories. Such overheads are reflected in the high price of the recombinant protein end-products.

To a lesser extent, animals have also been utilized for the production of recombinant proteins. Although large amounts of protein can be produced and relatively easily recovered from such animals (e.g., proteins specifically produced in mammary glands and secreted with the milk), production in such host is limited to the expression of proteins which do not interfere with the host physiology. In addition, transgenic animals are subject to lengthy lead times to develop herds with stable genetics, high operating costs, contamination by animal viruses and a relatively slow rate of biomass generation substantially prolonging the time period following which recovery of commercial amounts of the protein can be effected.

The biochemical, technical and economic limitations on existing prokaryotic and eukaryotic expression systems has created substantial

interest in developing new expression systems for the production of recombinant proteins.

Plants represent the most likely alternative to existing expression systems. With the availability and on going development of plant transformation techniques, most commercially important plant species can now be genetically modified to express a variety of recombinant proteins.

Such transformation techniques include, for example, the *Agrobacterium* vector system, which involves infection of the plant tissue with a bacterium (*Agrobacterium*) into which the foreign gene has been inserted. A number of methods for transforming plant cells with *Agrobacterium* are well known (Klee *et al.*, Annu. Rev. Plant Physiol. (1987) 38:467-486; Schell and Vasil Academic Publishers, San Diego, Calif. (1989) p. 2-25; and Gatenby (1989) in Plant Biotechnology, eds. Kung, S. and Arntzen, C. J., Butterworth Publishers, Boston, Mass. p. 93-112).

The biolistic or particle gun method, which permits genetic material to be delivered directly into intact cells or tissues by bombarding regeneratable tissues, such as meristems or embryogenic callus, with DNA-coated microparticles has contributed to plant transformation simplicity and efficiency. The microparticles penetrate the plant cells and act as inert carriers of a genetic material to be introduced therein. Microprojectile bombardment of embryogenic suspension cultures has proven successful for the production of transgenic plants of a variety of species. Various parameters that influence DNA delivery by particle bombardment have been defined (Klein *et al.*, Bio/Technology (1998) 6:559-563; McCabe *et al.*, Bio/Technology (1998) 6:923-926; and Sanford, Physiol. Plant. (1990) 79:206-209).

Micropipette systems are also used for the delivery of foreign DNA into plants via microinjection (Neuhaus *et al.*, Theor. Appl. Genet. (1987)

75:30-36; and Neuhaus and Spangenberg, *Physiol. Plant.* (1990) 79:213-217).

Other techniques developed to introduce foreign genes into plants include direct DNA uptake by plant tissue, or plant cell protoplasts (Schiell and Vasil (1987) Academic Publishers, San Diego, Calif. p. 52-68; and Toriyama *et al.*, *Bio/Technology* (1988) 6:1072-1074) or by germinating pollen (Chapman, Mantell and Daniels (1985) W. Longman, London, p. 197-209; and Ohta, *Proc. Natl. Acad. Sci. USA* (1986) 83:715-719).

DNA uptake induced by brief electric shock of plant cells has also been described (Zhang *et al.*, *Plant. Cell. Rep.* (1988) 7:379-384 and Fromm *et al.*, *Nature* (1986) 319:791-793).

In addition, virus mediated plant transformation has also been extensively described. Transformation of plants using plant viruses is described, for example, in U.S. Pat. No. 4,855,237 (BGV), EP-A 67,553 (TMV), Japanese Published Application No. 63-14693, EPA 194,809, EPA 278,667, and Gluzman *et al.*, (1988) *Communications in Molecular Biology: Viral Vectors*, Cold Spring Harbor Laboratory, New York, pp. 172-189. Pseudovirus particles for use in expressing foreign DNA in many hosts, including plants, have also been described, for reference, see, for example WO 87/06261.

The production of recombinant proteins and peptides in plants has been investigated using a variety of approaches including transcriptional fusions using a strong constitutive plant promoter (e.g., from cauliflower mosaic virus, Sijmons *et al.*, *Bio/Technology* (1990) 8:217-221); transcriptional fusions with organ specific promoter sequences (Radke *et al.*, *Theoret. Appl. Genet.* (1988) 75:685-694); and translational fusions which require subsequent cleavage of a recombinant protein (Vanderkerckove *et al.*, *Bio/Technology* (1989) 7:929-932).

The application of such genetic transformation techniques has allowed the incorporation of a variety of important genetic traits for crop improvement and also for the biotechnological production of extractable, valuable, foreign proteins including enzymes, vaccine proteins and antibodies.

Foreign proteins that have been successfully expressed in plant cells include proteins from bacteria (Fraley *et al.* Proc. Natl. Acad. Sci. U.S.A (1993) 80:4803-4807), animals (Misra and Gedamu, Theor. Appl. Genet. (1989) 78:161-168), fungi and other plant species (Fraley *et al.* Proc. Natl. Acad. Sci. U.S.A. (1983) 80:4803-4807). Some proteins, predominantly markers of DNA integration, have been expressed in specific cells and tissues including seeds (Sen Gupta-Gopalan *et al.* Proc. Natl. Acad. Sci. U.S.A. (1985) 82:3320-3324; Radke *et al.* Theor. Appl. Genet. (1988) 75:685-694).

Due to the advantageous economics of field-grown crops, the ability to synthesize proteins in storage organs like tubers, seeds, fruits and leaves and the ability of plants to perform many of the post-translational modifications previously described, several plant expression systems are currently investigated for potential as highly effective and economically feasible systems for the production of recombinant proteins.

Since highly expressive systems such as the ubiquitin fusion system described in U.S. Pat. No. 5,773,705 have been demonstrated, a major hurdle to an effective plant expression system resides with the relatively complicated purification procedures necessary in order to purify the recombinant protein.

As such, alternative expression approaches have been undertaken in an effort to simplify the purification procedure of the recombinant protein from the plant cells.

One such system focuses on the use of seed-storage protein promoters as a means of deriving seed-specific expression. Using such a system, Vanderkerckove *et al.*, (Bio/Technol. (1989) 7:929-932) expressed the peptide Leu-enkephalin in seeds of *Arabidopsis thaliana* and *Brassica napus*. The level of expression of this peptide was quite low and it appeared that expression of this peptide was limited to endosperm tissue.

Another system utilizing seeds as an expression host is disclosed in U.S. Pat. No. 5,888,789. This system provides for the secretion of heterologous protein by malting of monocot plant seeds. The heterologous genes are expressed during germination of the seeds and isolated from a malt.

U.S. Pat. No. 5,580,768 describes a method of producing a genetically transformed fluid-producing plant. The genetically transformed plant which can be for example, a rubber secreting (*Hevea*) plant is capable of expressing the target product in the fluid that it produces which in this case is latex.

U.S. Pat No. 5,650,554 describes the use of a class of genes called oil body protein genes, that have unique features, allowing the production of recombinant proteins that can be easily separated from other host cell components.

Many additional expression systems have been described utilizing specific targeting or directing of recombinant proteins to specific plant tissues.

Although systems which target or direct recombinant protein production to specific tissues allow for easier recombinant protein isolation such systems are typically limited in the effective host range and/or the amounts of recombinant proteins produced since such systems fail to exploit the entire plant biomass.

A novel approach for simplifying the purification of recombinant enzymes from plant host cells is disclosed in U.S. Pat. No. 5,474,925 which describes an expression construct utilizing a signal peptide translationally fused to a recombinant enzyme which targets the enzyme to the cellulose matrix of the cell wall. This enables the isolation of the enzyme along with the easily recoverable cellulose matrix. This system is utilized for the localized expression of commercially important enzymes in cotton fibers. According to this system, the expressed enzymes are recovered along with the cellulosic matter of the fibers. The enzyme-cellulose matrix recovered, is directly utilized for commercial enzymatic processes.

Although this system presents a simple means with which a recombinant protein can be expressed and isolated, it is limited to the production of enzymes in cotton fibers of the cotton plant.

Furthermore, a major hurdle encountered when expressing cellulose targeted proteins within a plant is the interference of the expressed products in the natural formation of the cell wall, which typically results in growth arrest of the plant growth. Although this hurdle can be overcome by, for example, targeting the protein to specific plant tissue as is the case for U.S. Pat. No. 5,474,925, this targeting severely limits the expressing biomass and as such the quantity of the expressed protein. In addition, targeting the expression to a specific plant tissue also limits the number of plant species which can be effectively utilized for such an expression.

There is thus a widely recognized need for, and it would be highly advantageous to have, a plant expression system and method which provide high level of expression of a recombinant protein and which allow simple and effective recovery of the expressed recombinant protein devoid of the above limitations.

SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a process of expressing a recombinant protein in a plant and of isolating the recombinant protein from the plant, the process comprising the steps of (a) providing a plant, a plant derived tissue or cultured plant cells expressing a fusion protein including the recombinant protein and a cellulose binding peptide being fused thereto, the fusion protein being compartmentalized within cells of the plant, plant derived tissue or cultured plant cells, so as to be sequestered from cell walls of the cells of the plant, plant derived tissue or cultured plant cells; (b) homogenizing the plant, plant derived tissue or cultured plant cells, so as to bring into contact the fusion protein with a cellulosic matter of the plant, plant derived tissue or cultured plant cells, to thereby effect affinity binding of the fusion protein via the cellulose binding peptide to the cellulosic matter, thereby obtaining a fusion protein cellulosic matter complex; and (c) isolating the fusion protein cellulosic matter complex.

According to further features in preferred embodiments of the invention described below, the process further comprising the steps of washing the fusion protein cellulosic matter complex, thereby removing endogenous plant proteins and other plant material therefrom and collecting the fusion protein cellulosic matter complex as a final product of the process.

According to still further features in the described preferred embodiments the process further comprising the steps of washing the fusion protein cellulosic matter complex, thereby removing endogenous plant proteins and other plant material therefrom, exposing the fusion protein cellulosic matter complex to conditions effective in dissociating the fusion protein from the cellulosic matter; and isolating the fusion protein, thereby obtaining an isolated fusion protein.

According to still further features in the described preferred embodiments the process further comprising the steps of exposing the isolated fusion protein to conditions effective in digesting the fusion protein so as to release the recombinant protein therefrom, thereby obtaining a released recombinant protein and isolating the released recombinant protein.

According to still further features in the described preferred embodiments the process further comprising the steps of washing the fusion protein cellulosic matter complex, thereby removing endogenous plant proteins and other plant material therefrom, exposing the fusion protein cellulosic matter complex to conditions effective in digesting the fusion protein so as to release the recombinant protein therefrom, thereby obtaining a released recombinant protein, and isolating the released recombinant protein.

According to still further features in the described preferred embodiments, the conditions effective in dissociating the fusion protein from the cellulosic matter are selected from the group consisting of basic conditions, denaturative conditions and affinity displacement conditions.

According to still further features in the described preferred embodiments, the conditions effective in digesting the fusion protein so as to release the recombinant protein therefrom are selected from the group consisting of proteolysis effected via a protease and proteolysis effected under predetermined cis or trans conditions for digesting a controllable intervening protein sequence.

According to another aspect of the present invention there is provided a genetically modified or viral infected plant or cultured plant cells expressing a fusion protein including a recombinant protein and a cellulose binding peptide

According to further features in preferred embodiments of the invention described below, the fusion protein is compartmentalized within

cells of the plant or cultured plant cells, so as to be sequestered from cell walls of the cells of the plant or cultured plant cells.

According to still further features in the described preferred embodiments the fusion protein is compartmentalized within a cellular compartment selected from the group consisting of cytoplasm, endoplasmic reticulum, golgi apparatus, oil bodies, starch bodies, chloroplastids, chloroplasts, chromoplastids, chromoplasts, vacuole, lysosomes, mitochondria, and nucleus.

According to still further features in the described preferred embodiments expression of the fusion protein is under a control of a constitutive or tissue specific plant promoter.

According to still further features in the described preferred embodiments the fusion protein includes a recombinant protein and a cellulose binding peptide separated therebetween via a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence.

According to yet another aspect of the present invention there is provided a composition of matter comprising (a) a plant derived cellulosic matter; and (b) a fusion protein including a recombinant protein and a cellulose binding peptide separated therebetween via a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence, the fusion protein being complexed to the plant derived cellulosic matter by affinity binding via the cellulose binding peptide.

According to still another aspect of the present invention there is provided a nucleic acid molecule comprising (a) a promoter sequence for directing protein expression in plant cells; (b) a heterologous nucleic acid sequence including (i) a first sequence encoding a cellulose binding peptide;

(ii) a second sequence encoding a recombinant protein, wherein the first and second sequences are joined together in frame; optionally (iii) a third sequence encoding a signal peptide for directing a protein to a cellular compartment, the third sequence being upstream and in frame with the first and second sequences; and/or optionally (iv) a fourth sequence encoding a
 5 unique amino acid sequence being recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence, the fourth sequence being between and in frame with the first and second sequences, wherein, the heterologous
 10 nucleic acid sequence being down stream the promoter sequence, such that expression of the heterologous nucleic acid sequence is effectable by the promoter sequence.

According to further features in preferred embodiments of the invention described below, the nucleic acid molecule further comprising a
 15 sequence element selected from the group consisting of an origin of replication for propagation in bacterial cells, at least one sequence element for integration into a plant's genome, a polyadenylation recognition sequence, a transcription termination signal, a sequence encoding a translation start site, a sequence encoding a translation stop site, plant RNA
 20 virus derived sequences, plant DNA virus derived sequences, tumor inducing (Ti) plasmid derived sequences, and a transposable element derived sequence.

The present invention successfully addresses the shortcomings of the presently known configurations by providing a novel process of expressing
 25 and isolating recombinant proteins and recombinant protein products from plants, plant derived tissue or cultured plant cells, which process exploits (i) the high affinity between cellulose binding peptides and cellulose; (ii) the inherent abundance of cellulose in *planta*; and (iii) the simplicity associated

with cellulose isolation from plants, plant derived tissue and/or cultured plant cells.

5

BRIEF DESCRIPTION OF THE DRAWINGS

The invention is herein described, by way of example only, with reference to the accompanying drawings, wherein:

10 FIG. 1 is a process flow chart demonstrating the process according to the present invention.

15 DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of a process which can be used for expressing and isolating recombinant proteins and recombinant protein products from plants, plant derived tissue or cultured plant cells. The present invention is further of nucleic acid molecules and genetically
20 modified or viral infected plants or plant cells which are useful while implementing the process, and further of a novel composition of matter which results from the process. Specifically, the present invention can be used to obtain large quantities of the recombinant proteins and the recombinant protein products in a simple and cost effective manner, since
25 the process according to the present invention exploits (i) the high affinity between cellulose binding peptides and cellulose; (ii) the inherent abundance of cellulose in *planta*; and (iii) the simplicity associated with cellulose isolation from plants, plant derived tissue and/or cultured plant cells.

The principles and operation of a process according to the present invention may be better understood with reference to the drawings and accompanying descriptions.

Before explaining at least one embodiment of the invention in detail,
5 it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology
10 employed herein is for the purpose of description and should not be regarded as limiting.

Process:

Referring now to the drawings, Figure 1 is a flow chart illustrating the process according to the teachings of the present invention.

15 The process according to the present invention is directed at expressing a recombinant protein in a plant and further at isolating the recombinant protein from the plant. The process according to the present invention is effected by first providing a plant, a plant derived tissue or cultured plant cells (which are referred to herein below individually and
20 collectively as "plant material") 10 expressing a fusion protein which includes the recombinant protein and a cellulose binding peptide fused thereto. The fusion protein is compartmentalized within cells of the plant material, so as to be sequestered from cell walls of the cells of the plant material. As used herein in the specification and in the claims section that
25 follows, the phrase "cultured plant cells" includes both non-differentiated plant cell cultures and some what more differentiated callus cultures.

Compartmentalizing and thereby sequestering the fusion protein from the cell walls is an essential feature of the present invention because high levels of expressed cellulose binding peptide associated with plant

cell walls inhibit plant growth to a great extent. See to this effect U.S. Pat. applications Nos. 09/006,632; 09/006,636; and PCT/IL98/00345 (WO 99/07830).

When sufficient expression has been detected by sampling and testing the plant material as further detailed hereinunder, the plant material is homogenized 12 so as to bring into contact the fusion protein with a cellulosic matter of the plant material, to thereby effect affinity binding 14 of the fusion protein via the cellulose binding peptide to the cellulosic matter, thereby obtaining a fusion protein cellulosic matter complex. Conditions such as, but not limited to, temperature, pH, salt concentration, time and the like are preferably set so as to allow maximal binding. Such conditions are well known to the skilled artisan and can be experimentally modified to best suit a specific application. Sampling and testing can be employed to monitor the binding process, as further detailed hereinunder.

When sufficient binding has occurred the fusion protein cellulosic matter complex is collected or isolated 16 by methods well known to the skilled artisan which methods are traditionally employed for isolation of cellulosic matter from plant material. Thereafter, a wash step 18 is employed to remove unbound material, including, in particular, unbound endogenous plant proteins, thereby isolating the fusion protein cellulosic matter complex. The wash step can be repeated one or several times with a single or several wash solutions, each of which can include in addition to water, buffers, salts, detergents and the like to efficiently effect the removal of unbound matter from the fusion protein cellulosic matter complex. The wash step can be effected in solution using appropriate stirring, however, advantageously, the wash step is effected within a column into which the collected or isolated fusion protein cellulosic matter complex is packed and subsequently washed.

According to one embodiment of the present invention, and as indicated in Figure 1 by numeral 20, the fusion protein cellulosic matter complex is collected as a final product of the process. Such a final product can serve as a pack for affinity columns. In this case the recombinant protein is selected to have affinity to a ligand, which can then be affinity purified via a column packed with the fusion protein cellulosic matter complex, in a manner otherwise similar to that described in U.S. Pat. No. 5,474,925, which is incorporated herein by reference. One of the advantages of the process described herein over the teachings of U.S. Pat. No. 5,474,925 is that by sequestering the fusion protein from the cell walls one can achieve very high expression of the fusion protein as compared to the low expression levels practically enabled by U.S. Pat. No. 5,474,925, because no deleterious effect on plant growth is exerted. As a result, the specific activity of the fusion protein cellulosic matter complex formed according to the present invention, i.e., the number of fusion protein molecules per weight of cellulosic matter, is far superior. Further details relating to the effect of high cellulose binding peptide expression on plant development see also PCT/IL98/000345.

According to another embodiment of the present invention, as indicated by numeral 22, the final product of the process according to the present invention is the fusion protein itself 24. Thus, according to this embodiment of the present invention, conditions effective in dissociating the fusion protein from the cellulosic matter are used to effect such dissociation. The dissociated fusion protein is thereafter readily isolated by any conventional separation technique, such as, but not limited to, elution or size separation, such as differential filtration or centrifugation, thereby obtaining an isolated fusion protein. Conditions effective in dissociating the fusion protein from the cellulosic matter include, but are not limited to, basic conditions (e.g., 20 mM Tris pH 12) which are known to dissociate all

cellulose binding peptides from cellulose, denaturative conditions, or affinity displacement conditions, e.g., using 200 nM glucose or cellobiose which are known in their ability to elute family IX cellulose binding domains (CBDs). Alternatively, a protein cleavage site can be inserted in the cellulose binding peptide to facilitate the dissociation of the fusion protein by specific proteolysis, for example. See to this effect and to other uses of CBD-fusion proteins U.S. Pat. Nos. 5,719,044; 5,670,623; 5,856,021; 5,137,819; 5,202,247; 5,340,731; and 5,474,925; and U.S. Pat. applications Nos. 08/788,621; and 08/788,622; EP 0 381 719 B1, and EP application No. 93907724.4. See also the teachings of U.S. Pat. No. 5,834,247, which is further described hereinunder.

As indicated by numeral 26, the fusion protein thus isolated can be exposed to conditions effective in digesting the fusion protein so as to release the recombinant protein therefrom, thereby obtaining a released recombinant protein which can be thereafter isolated as a final product 28. Conditions effective in digesting the fusion protein so as to release the recombinant protein therefrom include, but are not limited to, proteolysis effected via a protease, such as, but not limited to, Factor Xa, enterokinase, thrombin, trypsin, papain, pepsin, chymotrypsin and the like, or proteolysis effected via controllable intervening protein sequence (CIVPS) inserted into or adjacent the cellulose binding peptide, the CIVPS are capable of excision from or cleavage of the peptide under predetermined conditions in cis or in trans, e.g., increase in temperature, exposure to light, unblocking of amino acid residues by dephosphorylation and treatment with chemical reagents or deglycosylation, examples include proteolysis effected under acidic conditions (HCl, e.g., to cleave between Asp and Pro) and proteolysis effected by a proteolyzing reagent, such as CNBr to cleave downstream of Met, all as known in the art and/or as further described in U.S. Pat. No. 5,834,247, which is incorporated herein by reference.

Isolating final product 28 from other proteolytic products derived, for example, from the cellulose binding peptide, can be effected by any one of a number of protein isolation techniques well known to the skilled artisan, including, but not limited to, affinity separation via, for example, antibodies
5 bound to a solid support, size and/or charge based separation via gel electrophoresis or chromatography, and the like. Additional methods include, but are not limited to, fractionation, gel-filtration, ion-exchange, hydrophobic, and affinity chromatography, ultrafiltration and crystallization.

10 According to an alternative embodiment of the process of the present invention, as indicated in Figure 1 by numeral 30, the washed fusion protein cellulose matter complex resulting from step 18 is exposed to conditions effective in digesting the fusion protein so as to release the recombinant protein therefrom, thereby obtaining a released recombinant protein. These
15 conditions are similar to those described with respect to step 26. The released recombinant protein is thereafter readily isolated by any conventional separation technique, such as, but not limited to, displacement or size separation, such as differential filtration or centrifugation, thereby obtaining an isolated recombinant protein final product 32.

20 ***Cellulose binding peptides:***

As used herein in the specification and in the claims section below, the phrase "cellulose binding peptide" includes peptides e.g., proteins and domains (portions) thereof, which are capable of, when expressed in plant cells, affinity binding to a plant derived cellulosic matter following
25 homogenization and cell rupture. The phrase thus includes, for example, peptides which were screened for their cellulose binding activity out of a library, such as a peptide library or a DNA library (e.g., a cDNA expression library or a display library) and the genes encoding such peptides isolated and are expressible in plants. Yet, the phrase also

includes peptides designed and engineered to be capable of binding to cellulose and/or units thereof.

Such peptides include amino acid sequences expressible in plants that are originally derived from a cellulose binding region of, e.g., a cellulose binding protein (CBP) or a cellulose binding domain (CBD). The cellulose binding peptide according to the present invention can include any amino acid sequence expressible in plants which binds to a cellulose polymer. For example, the cellulose binding domain or protein can be derived from a cellulase, a binding domain of a cellulose binding protein or a protein screened for, and isolated from, a peptide library, or a protein designed and engineered to be capable of binding to cellulose or to saccharide units thereof, and which is expressible in plants. The cellulose binding domain or protein can be naturally occurring or synthetic, as long as it is expressible in plants. Suitable polysaccharidases from which a cellulose binding domain or protein expressible in plants may be obtained include β -1,4-glucanases. In a preferred embodiment, a cellulose binding domain or protein from a cellulase or scaffoldin is used. Typically, the amino acid sequence of the cellulose binding peptide expressed in plants according to the present invention is essentially lacking in the hydrolytic activity of a polysaccharidase (e.g., cellulase, chitinase), but retains the cellulose binding activity. The amino acid sequence preferably has less than about 10 % of the hydrolytic activity of the native polysaccharidase; more preferably less than about 5 %, and most preferably less than about 1 % of the hydrolytic activity of the native polysaccharidase, ideally no activity altogether.

The cellulose binding domain or protein can be obtained from a variety of sources, including enzymes and other proteins which bind to cellulose which find use in the subject invention.

In Table 4 below are listed those binding domains which bind to one or more soluble/insoluble polysaccharides including all binding domains with affinity for soluble glucans (α , β , and/or mixed linkages). The N1 cellulose-binding domain from endoglucanase CenC of *C. fimi* is the only protein known to bind soluble cellosaccharides and one of a small set of proteins which are known to bind any soluble polysaccharides. Also, listed in Tables 1 to 3 are examples of proteins containing putative β -1,3-glucan-binding domains (Table 1); proteins containing Streptococcal glucan-binding repeats (Cpl superfamily) (Table 2); and enzymes with chitin-binding domains, which may also bind cellulose (Table 3). The genes encoding each one of the peptides listed in Tables 1-4 are either isolated or can be isolated as further detailed hereinunder, and therefore, such peptides are expressible in plants. Scaffoldin proteins or portions thereof, which include a cellulose binding domain, such as that produced by *Clostridium cellulovorans* (Shoseyov *et al.*, PCT/US94/04132) can also be used as the cellulose binding peptide expressible in plants according to the present invention. Several fungi, including *Trichoderma* species and others, also produce polysaccharidases from which polysaccharide binding domains or proteins expressible in plants can be isolated. Additional examples can be found in, for example, Microbial Hydrolysis of Polysaccharides, R. A. J. Warren, Annu. Rev. Microbiol. 1996, 50:183-212; and "Advances in Microbial Physiology" R. K. Poole, Ed., 1995, Academic Press Limited, both are incorporated by reference as if fully set forth herein.

Table 1
Overview of proteins containing putative β -1,3 glucan-binding domains

Source (strain)	Protein	accession No.	Ref ¹
Type I			
<i>B. circulans</i> (WL-12)	GLCA1	P23903/M34503/JQ0420	1
<i>B. circulans</i> (IAM 1165)	BglH	JN0772/D17519/S67033	2
Type II			
<i>Actinomadura</i> sp. (FC7)	XynII	U08894	3
<i>Arthrobacter</i> sp. (YCWD3)	GLCI	D23668	9
<i>O. xanthineolytica</i>	GLC	P22222/M60826/A39094	4
<i>R. faecitabidus</i> (YLM-50)	RP I	Q05308/A45053/D10753	5a,b
<i>R. communis</i>		A12892	6
<i>S. lividans</i> (1326)	XlnA	P26514/M64551/JS07986	7
<i>T. tridentatus</i>	FactorGa	D16622	8

B. : *Bacillus*, *O.* : *Oerskovia*, *R. faecitabidus* : *Rarobacter faecitabidus*, *R. communis*: *Ricinus communis*, *S.* : *Streptomyces*, *T.* : *Tachypleus* (Horseshoe Crab)

¹References:

- 1) Yahata *et al.* (1990) *Gene* 86, 113-117
- 2) Yamamoto *et al.* (1993) *Biosci. Biotechnol. Biochem.* 57, 1518-1525
- 3) Harpin *et al.* (1994) *EMBL Data Library*
- 4) Shen *et al.* (1991) *J. Biol. Chem.* 266, 1058-1063
- 5a) Shimoi *et al.* (1992) *J. Biol. Chem.* 267, 25189-25195
- 5b) Shimoi *et al.* (1992) *J. Biochem* 110, 608-613
- 6) Horn *et al.* (1989) Patent A12892
- 7) Shareck *et al.* (1991) *Gene* 107, 75-82
- 8) Seki *et al.* (1994) *J. Biol. Chem.* 269, 1370-1374
- 9) Watanabe *et al.* (1993) *EMBL Data Library*

Table 2
Overview of proteins containing *Streptococcal* glucan-binding repeats (*Cpl* superfamily)

Source	Protein	Accession No.	Ref. ²
<i>S. downei</i> (<i>sobrinus</i>) (0MZ176)	GTF-I	D13858	1
<i>S. downei</i> (<i>sobrinus</i>) (MFe28)	GTF-I	P11001/M17391	2
<i>S. downei</i> (<i>sobrinus</i>) (MFe28)	GTF-S	P29336/M30943/A41483	3
<i>S. downei</i> (<i>sobrinus</i>) (6715)	GTF-I	P27470/D90216/A38175	4
<i>S. downei</i> (<i>sobrinus</i>)	DEI	L34406	5
<i>S. mutants</i> (Ingbritt)	GBP	M30945/A37184	6
<i>S. mutants</i> (GS-5)	GTF-B	A33128	7
<i>S. mutants</i> (GS-5)	GTF-B	P08987/M17361/B33135	8
<i>S. mutants</i>	GTF-B ^{3'} -ORF	P05427/C33135	8

	<i>S. mutants</i> (GS-5)	GTF-C	P13470/M17361/M22054	9
	<i>S. mutants</i> (GS-5)	GTF-C	not available	10
	<i>S. mutants</i> (GS-5)	GTF-D	M29296/A45866	11
5	<i>S. salivarius</i>	GTF-J	A44811/S22726/S28809	12
			Z11873/M64111	
	<i>S. salivarius</i>	GTF-K	S22737/S22727/Z11872	13
	<i>S. salivarius</i> (ATCC25975)	GTF-L	L35495	14
	<i>S. salivarius</i> (ATCC25975)	GTF-M	L35928	14
10				
	<i>S. pneumoniae</i> R6	LytA	P06653/A25634/M13812	15
	<i>S. pneumoniae</i>	PspA	A41971/M74122	16
	Phage HB-3	HBL	P32762/M34652	17
15	Phage Cp-1	CPL-1	P15057/J03586/A31086	18
	Phage Cp-9	CPL-9	P19386/M34780/JQ0438	19
	Phage EJ-1	EJL	A42936	20
	<i>C. difficile</i> (VPI 10463)	ToxA	P16154/A37052/M30307	21
20			X51797/S08638	
	<i>C. difficile</i> (BARTS W1)	ToxA	A60991/X17194	22
	<i>C. difficile</i> (VPI 10463)	ToxB	P18177/X53138/X60984	23,24
			S10317	
	<i>C. difficile</i> (1470)	ToxB	S44271/Z23277	25,26
25				
	<i>C. novyi</i>	a-toxin	S44272/Z23280	27
	<i>C. novyi</i>	a-toxin	Z48636	28
	<i>C. acetobutylicum</i> (NCIB8052)	CspA	S49255/Z37723	29
30	<i>C. acetobutylicum</i> (NCIB8052)	CspB	Z50008	30
	<i>C. acetobutylicum</i> (NCIB8052)	CspC	Z50033	30
	<i>C. acetobutylicum</i> (NCIB8052)	CspD	Z50009	30

References:

- 1) Sato *et al.* (1993) *DNA sequence* 4, 19-27
- 2) Ferreti *et al.* (1987) *J. Bacteriol.* 169, 4271-4278
- 3) Gilmore *et al.* (1990) *J. Infect. Immun.* 58, 2452-2458
- 4) Abo *et al.* (1991) *J. Bacteriol.* 173, 989-996
- 5) Sun *et al.* (1994) *J. Bacteriol.* 176, 7213-7222
- 6) Banas *et al.* (1990) *J. Infect. Immun.* 58, 667-673
- 7) Shiroza *et al.* (1990) *Protein Sequence Database*
- 8) Shiroza *et al.* (1987) *J. Bacteriol.* 169, 4263-4270
- 9) Ueda *et al.* (1988) *Gene* 69, 101-109
- 10) Russel (1990) *Arch. Oral. Biol.* 35, 53-58
- 11) Honda *et al.* (1990) *J. Gen. Microbiol.* 136, 2099-2105
- 12) Giffard *et al.* (1991) *J. Gen. Microbiol.* 137, 2577-2593
- 13) Jacques (1992) *EMBL Data Library*
- 14) Simpson *et al.* (1995) *J. Infect. Immun.* 63, 609-621
- 15) Gargia *et al.* (1986) *Gene* 43, 265-272
- 16) Yother *et al.* (1992) *J. Bacteriol.* 174, 601-609
- 17) Romero *et al.* (1990) *J. Bacteriol.* 172, 5064-5070
- 18) Garcia *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85, 914-918
- 19) Garcia *et al.* (1990) *Gene* 86, 81-88
- 20) Diaz *et al.* (1992) *J. Bacteriol.* 174, 5516-5525
- 21) Dove *et al.* (1990) *J. Infect. Immun.* 58, 480-488
- 22) Wren *et al.* (1990) *FEMS Microbiol. Lett.* 70, 1-6

- 23) Barroso *et al.* (1990) *Nucleic Acids Res.* 18, 4004-4004
- 24) von Eichel-Streiber *et al.* (1992) *Mol. Gen. Genet.* 233, 260-268
- 25) Sarteringer *et al.* (1993) *EMBL Data Library*
- 26) von Eichel-Streiber *et al.* (1995) *Mol. Microbiol.* In Press
- 27) Hofmann *et al.* (1993) *EMBL Data Library*
- 28) Hofmann *et al.* (1995) *Mol. Gen. Genet.* In Press
- 29) Sanchez *et al.* (1994) *EMBL Data Library*
- 30) Sanchez *et al.* (1995) *EMBL Data Library*

New cellulose binding peptides with interesting binding characteristics and specificities can be identified and screened for and the genes encoding same isolated using well known molecular biology approaches combined with a variety of other procedures including, for example, spectroscopic (titration) methods such as: NMR spectroscopy (Zhu *et al.* *Biochemistry* (1995) 34:13196-13202, Gehring *et al.* *Biochemistry* (1991) 30:5524-5531), UV difference spectroscopy (Belshaw *et al.* *Eur. J. Biochem.* (1993) 211:717-724), fluorescence (titration) spectroscopy (Miller *et al.* *J. Biol. Chem.* (1983) 258:13665-13672), UV or fluorescence stopped flow analysis (De Boeck *et al.* *Eur. J. Biochem.* (1985) 149:141-415), affinity methods such as affinity electrophoresis (Mimura *et al.* *J. chromatography* (1992) 597:345-350) or affinity chromatography on immobilized mono or oligosaccharides, precipitation or agglutination analysis including turbidimetric or nephelometric analysis (Knibbs *et al.* *J. Biol. Chem.* (1993) 14940-14947), competitive inhibition assays (with or without quantitative IC50 determination) and various physical or physico-chemical methods including differential scanning or isothermal titration calorimetry (Sigurskjold *et al.* *J. Biol. Chem.* (1992) 267:8371-8376; Sigurskjold *et al.* *Eur. J. Biol.* (1994) 225:133-141) or comparative protein stability assays (melts) in the absence or presence of oligo saccharides using thermal CD or fluorescence spectroscopy.

The K_a for binding of the cellulose binding domains or proteins to cellulose is at least in the range of weak antibody-antigen extractions, i.e., $\geq 10^3$, preferably 10^4 , most preferably 10^6 M^{-1} . If the binding of the

cellulose binding domain or protein to cellulose is exothermic or endothermic, then binding will increase or decrease, respectively, at lower temperatures, providing a means for temperature modulation of the binding step, see numeral 14 in Figure 1.

5

Table 3
Overview of enzymes with chitin-binding domains

	Source (strain)	Enzyme	Accession No.	Ref. ³
10	<hr/>			
	Bacterial enzymes			
	Type I			
15	<i>Aeromonas</i> sp. (No10S-24)	Chi	D31818	1
	<i>Bacillus circulans</i> (WL-12)	ChiA1	P20533/M57601/A38368	2
	<i>Bacillus circulans</i> (WL-12)	ChiD	P27050/D10594	3
	<i>Janthinobacterium lividum</i>	Chi69	U07025	4
20	<i>Streptomyces griseus</i>	Protease C	A53669	5
	Type II			
	<i>Aeromonas cavia</i> (K1)	Chi	U09139	6
25	<i>Alteromonas</i> sp (0-7)	Chi85	A40633/P32823/D13762	7
	<i>Autographa californica</i> (C6)	NPH-128 ^a	P41684/L22858	8
	<i>Serratia marcescens</i>	ChiA	A25090/X03657/L01455/P07254	9
	Type III			
30	<i>Rhizopus oligosporus</i> (IFO8631)	Chi1	P29026/A47022/D10157/S27418	10
	<i>Rhizopus oligosporus</i> (IFO8631)	Chi2	P29027/B47022/D10158/S27419	10
	<i>Saccharomyces cerevisiae</i>	Chi	S50371/U17243	11
35	<i>Saccharomyces cerevisiae</i> (DBY939)	Chi1	P29028/M74069	12
	<i>Saccharomyces cerevisiae</i> (DBY918)	Chi2	P29029/M7407/B41035	12
40	Plant enzymes			
	Hevein superfamily			
	<i>Allium sativum</i>	Chi	M94105	13
45	<i>Amaranthus caudatus</i>	AMP-1 ^b	P27275/A40240	14, 15
	<i>Amaranthus caudatus</i>	AMP-2 ^b	S37381/A40240	14, 15
	<i>Arabidopsis thaliana</i> (cv. colombia)	ChiB	P19171/M38240/B45511	16
	<i>Arabidopsis thaliana</i>	PHP ^c	U01880	17
50	<i>Brassica napus</i>	Chi	U21848	18
	<i>Brassica napus</i>	Chi2	Q09023/M95835	19
	<i>Hevea brasiliensis</i>	Hev1 ^d	P02877/M36986/A03770/A38288	20, 21

	<i>Hordeum vulgare</i>	Chi33	L34211	22
	<i>Lycopersicon esculentum</i>	Chi9	Q05538/Z15140/S37344	23
	<i>Nicotiana tabacum</i>	CBP20 ^c	S72424	24
	<i>Nicotiana tabacum</i>	Chi	A21091	25
5	<i>Nicotiana tabacum</i> (cv. Havana)	Chi	A29074/M15173/S20981/S19855	26
	<i>Nicotiana tabacum</i> (FB7-1)	Chi	JQ0993/S0828	27
	<i>Nicotiana tabacum</i> (cv. Samsun)	Chi	A16119	28
	<i>Nicotiana tabacum</i> (cv. Havana)	Chi	P08252/X16939/S08627	27
	<i>Nicotiana tabacum</i> (cv. BY4)	Chi	P24091/X51599/X64519/S13322	26,27,29
10	<i>Nicotiana tabacum</i> (cv. Havana)	Chi	P29059/X64518/S20982	26
	<i>Oryza sativum</i> (IR36)	ChiA	L37289	30
	<i>Oryza sativum</i>	ChiB	JC2253/S42829/Z29962	31
	<i>Oryza sativum</i>	Chi	S39979/S40414/X56787	32
	<i>Oryza sativum</i> (cv. Japonicum)	Chi	X56063	33
15	<i>Oryza sativum</i> (cv. Japonicum)	Chi1	P24626/X54367/S14948	34
	<i>Oryza sativum</i>	Chi2	P25765/S15997	35
	<i>Oryza sativum</i> (cv. Japonicum)	Chi3	D16223	
	<i>Oryza sativum</i>	ChiA	JC2252/S42828	30
	<i>Oryza sativum</i>	Chi1	D16221	32
20	<i>Oryza sativum</i> (IR58)	Chi	U02286	36
	<i>Oryza sativum</i>	Chi	X87109	37
	<i>Pisum sativum</i> (cv. Birte)	Chi	P36907/X63899	38
	<i>Pisum sativum</i> (cv. Alcan)	Chi2	L37876	39
	<i>Populus trichocarpa</i>	Chi	S18750/S18751/X59995/P29032	40
25	<i>Populus trichocarpa</i> (H11-11)	Chi	U01660	41
	<i>Phaseolus vulgaris</i> (cv. Saxa)	Chi	A24215/S43926/Jq0965/P36361	42
	<i>Phaseolus vulgaris</i> (cv. Saxa)	Chi	P06215/M13968/M19052/A25898	43,44,45
	<i>Sambucus nigra</i>	PR-3 ^f	Z46948	46
	<i>Secale cereale</i>	Chi	JC2071	47
30	<i>Solanum tuberosum</i>	ChiB1	U02605	48
	<i>Solanum tuberosum</i>	ChiB2	U02606	48
	<i>Solanum tuberosum</i>	ChiB3	U02607/S43317	48
	<i>Solanum tuberosum</i>	ChiB4	U02608	48
	<i>Solanum tuberosum</i>	WIN-1 ^g	P09761/X13497/S04926	49
35	(cv. Maris Piper)			
	<i>Solanum tuberosum</i>	WIN-2 ^g	P09762/X13497/S04927	49
	(cv. Maris Piper)			
	<i>Triticum aestivum</i>	Chi	S38670/X76041	50
	<i>Triticum aestivum</i>	WGA-1 ^h	P10968/M25536/S09623/S07289	51,52
40	<i>Triticum aestivum</i>	WGA-2 ^h	P02876/M25537/S09624	51,53
	<i>Triticum aestivum</i>	WGA-3	P10969/I02961/S10045/A28401	54
	<i>Ulmus americana</i> (NPS3-487)	Chi	L22032	55
	<i>Urtica dioica</i>	AGL ⁱ	M87302	56
	<i>Vigna unguiculata</i>	Chi1	X88800	57
45	(cv. Red caloonna)			

^aNHP : nuclear polyhedrosis virus endochitinase like sequence; Chi : chitinase, ^banti-microbial peptide, ^cpre-hevein like protein, ^dhevein, ^echitin-binding protein, ^fpathogenesis related protein, ^gwound-induced protein, ^hwheat germ agglutinin, ⁱagglutinin (lectin).

³References:

- 1) Udca *et al.* (1994) *J. Ferment. Bioeng.* 78, 205-211
- 2) Watanabe *et al.* (1990) *J. Biol. Chem.* 265, 15659-16565
- 3) Watanabe *et al.* (1992) *J. Bacteriol.* 174, 408-414
- 4) Gleave *et al.* (1994) *EMBL Data Library*
- 5) Sidhu *et al.* (1994) *J. Biol. Chem.* 269, 20167-20171
- 6) Jones *et al.* (1986) *EMBO J.* 5, 467-473

- 7) Sitrit *et al.* (1994) *EMBL Data Library*
- 8) Genbank entry only
- 9) Tsujibo *et al.* (1993) *J. Bacteriol.* **175**, 176-181
- 10) Yanai *et al.* (1992) *J. Bacteriol.* **174**, 7398-7406
- 5 11) Pauley (1994) *EMBL Data Library*
- 12) Kuranda *et al.* (1991) *J. Biol. Chem.* **266**, 19758-19767
- 13) van Damme *et al.* (1992) *EMBL Data Library*
- 14) Broekaert *et al.* (1992) *Biochemistry* **31**, 4308-4314
- 15) de Bolle *et al.* (1993) *Plant Mol. Physiol.* **22**, 1187-1190
- 10 16) Samac *et al.* (1990) *Plant Physiol.* **93**, 907-914
- 17) Potter *et al.* (1993) *Mol. Plant Microbe Interact.* **6**, 680-685
- 18) Buchanan-Wollaston (1995) *EMBL Data Library*
- 19) Hamel *et al.* (1993) *Plant Physiol.* **101**, 1403-1403
- 20) Broekaert *et al.* (1990) *Proc. Natl. Acad. Sci. USA* **87**, 7633-7637
- 15 21) Lee *et al.* (1991) *J. Biol. Chem.* **266**, 15944-15948
- 22) Leah *et al.* (1994) *Plant Physiol.* **6**, 579-589
- 23) Danhash *et al.* (1993) *Plant Mol. Biol.* **22** 1017-1029
- 24) Ponstein *et al.* (1994) *Plant Physiol.* **104**, 109-118
- 25) Meins *et al.* (1991) Patent EP0418695-A1
- 20 26) van Buuren *et al.* (1992) *Mol. Gen. Genet.* **232**, 460-469
- 27) Shinshi *et al.* (1990) *Plant Mol. Biol.* **14**, 357-368
- 28) Cornellisen *et al.* (1991) Patent EP0440304-A2
- 29) Fukuda *et al.* (1991) *Plant Mol. Biol.* **16**, 1-10
- 30) Yun *et al.* (1994) *EMBL Data Library*
- 25 31) Kim *et al.* (1994) *Biosci. Biotechnol. Biochem.* **58**, 1164-1166
- 32) Nishizawa *et al.* (1993) *Mol. Gen. Genet.* **241**, 1-10
- 33) Nishizawa *et al.* (1991) *Plant Sci* **76**, 211-218
- 34) Huang *et al.* (1991) *Plant Mol. Biol.* **16**, 479-480
- 35) Zhu *et al.* (1991) *Mol. Gen. Genet.* **226**, 289-296
- 30 36) Muthukrishnan *et al.* (1993) *EMBL Data Library*
- 37) Xu (1995) *EMBL Data Library*
- 38) Vad *et al.* (1993) *Plant Sci* **92**, 69-79
- 39) Chang *et al.* (1994) *EMBL Data Library*
- 40) Davis *et al.* (1991) *Plant Mol. Biol.* **17**, 631-639
- 35 41) Clarke *et al.* (1994) *Plant Mol. Biol.* **25**, 799-815
- 42) Broglie *et al.* (1989) *Plant Cell* **1**, 599-607
- 43) Broglie *et al.* (1986) *Proc. Natl. acad. Sci. USA* **83**, 6820-6824
- 44) Lucas *et al.* (1985) *FEBS Lett.* **193**, 208-210
- 45) Hedrick *et al.* (1988) *Plant Physiol.* **86**, 182-186
- 40 46) Roberts *et al.* (1994) *EMBL Data Library*
- 47) Vamagami *et al.* (1994) *Biosci. Biotechnol. Biochem.* **58**, 322-329
- 48) Beerhues *et al.* (1994) *Plant Mol. Biol.* **24**, 353-367
- 49) Stanford *et al.* (1989) *Mol. Gen. Genet.* **215**, 200-208
- 50) Liao *et al.* (1993) *EMBL Data Library*
- 45 51) Smith *et al.* (1989) *Plant Mol. Biol.* **13**, 601-603
- 52) Wright *et al.* (1989) *J. Mol. Evol.* **28**, 327-336
- 53) Wright *et al.* (1984) *Biochemistry* **23**, 280-287
- 54) Raikhel *et al.* (1987) *Proc. Natl. acad. Sci. USA* **84**, 6745-6749
- 55) Hajela *et al.* (1993) *EMBL Data Library*
- 50 56) Lerner *et al.* (1992) *J. Biol. Chem.* **267**, 11085-11091
- 57) Vo *et al.* (1995) *EMBL Data Library*

Table 4
Sources of polysaccharide binding domains

	Binding Domain	Proteins Where Binding Domain is Found
5		
10	Cellulose Binding Domains ¹	β -glucanases (avicelases, CMCase, cellodextrinases) exoglucanases or cellobiohydrolases cellulose binding proteins xylanases mixed xylanases/glucanases
15		esterases chitinases β -1,3-glucanases β -1,3-(β -1,4)-glucanases (β -)mannanases
20		β -glucosidases/galactosidases cellulose synthases (unconfirmed)
25	Starch/Maltodextrin Binding Domains	α -amylases ^{2,3} β -amylases ^{4,5} pullulanases glucoamylases ^{6,7} cyclodextrin glucotransferases ⁸⁻¹⁰ (cyclomaltodextrin glucanotransferases) maltodextrin binding proteins ¹¹
30	Dextran Binding Domains	(<i>Streptococcal</i>) glycosyl transferases ¹² dextran sucrases (unconfirmed) <i>Clostridial</i> toxins ^{13,14} glucoamylases ⁶ dextran binding proteins
35	β -Glucan Binding Domains	β -1,3-glucanases ^{15,16} β -1,3-(β -1,4)-glucanases (unconfirmed) β -1,3-glucan binding protein
40	Chitin Binding Domains	chitinases chitobiases chitin binding proteins (see also cellulose binding domains)
45		Hevein

¹Gilkes *et al.*, *Adv. Microbiol. Reviews*, (1991) 303-315.

²Sogaard *et al.*, *J. Biol. Chem.* (1993) 268:22480.

³Weselake *et al.*, *Cereal Chem.* (1983) 60:98.

⁴Svensson *et al.*, *J.* (1989) 264:309.

⁵Jespersen *et al.*, *J.* (1991) 280:51.

⁶Belshaw *et al.*, *Eur. J. Biochem.* (1993) 211:717.

⁷Sigurskjold *et al.*, *Eur. J. Biochem.* (1994) 225:133.

⁸Villette *et al.*, *Biotechnol. Appl. Biochem.* (1992) 16:57.

⁹Fukada *et al.*, *Biosci. Biotechnol. Biochem.* (1992) 56:556.

¹⁰Lawson *et al.*, *J. Mol. Biol.* (1994) 236:590.

50

55

¹⁴von Eichel-Streiber *et al.*, *Mol. Gen. Genet.* (1992) 233:260.

¹⁵Klebl *et al.*, *J. Bacteriol.* (1989) 171:6259.

¹⁶Watanabe *et al.*, *J. Bacteriol.* (1992) 174:186.

¹⁷Duvic *et al.*, *J. Biol. Chem.* (1990) :9327.

5 Thus, and as already stated, the phrase "polysaccharide binding peptide" includes an amino acid sequence which comprises at least a functional portion of a polysaccharide binding region (domain) of a polysaccharidase or a polysaccharide binding protein. The phrase further relates to a polypeptide screened for its cellulose binding activity out of a
10 library, such as a peptide library or a DNA library (e.g., a cDNA library or a display library). By "functional portion" is intended an amino acid sequence which binds to cellulose.

 The techniques used in isolating polysaccharidase genes, such as cellulase genes, and genes for cellulose binding proteins are known in the
15 art, including synthesis, isolation from genomic DNA, preparation from cDNA, or combinations thereof. (See, U.S. Pat. Nos. 5,137,819; 5,202,247; 5,340,731; 5,496,934; and 5,837,814). The sequences for several binding domains, which bind to soluble oligosaccharides are known (See, Figure 1 of PCT/CA97/00033, WO 97/26358). The DNAs
20 coding for a variety of polysaccharidases and polysaccharide binding proteins are also known. Various techniques for manipulation of genes are well known, and include restriction, digestion, resection, ligation, *in vitro* mutagenesis, primer repair, employing linkers and adapters, and the like (see Sambrook *et al.*, *Molecular Cloning - A Laboratory Manual*,
25 Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989, which is incorporated herein by reference).

 The amino acid sequence of a polysaccharidase can be used to design a probe to screen a cDNA or a genomic library prepared from mRNA or DNA from cells of interest as donor cells for a
30 polysaccharidase gene or a polysaccharide binding protein gene. By using the polysaccharidase cDNA or binding protein cDNA or a fragment

thereof as a hybridization probe, structurally related genes found in other species can be easily cloned and provide a cellulose binding peptide which is expressible in plants according to the present invention. Particularly contemplated is the isolation of genes from organisms that express
 5 polysaccharidase activity using oligonucleotide probes based on the nucleotide sequences of genes obtainable from an organism wherein the catalytic and binding domains of the polysaccharidase are discrete, although other polysaccharide binding proteins also can be used (see, for example, Shoseyov, *et al.*, Proc. Nat'l. Acad. Sci. (USA) (1992)
 10 89:3483-3487).

Probes developed using consensus sequences for the binding domain of a polysaccharidase or polysaccharide-binding protein are of particular interest. The β -1,4-glycanases from *C. fimi* characterized to date are endoglucanases A, B, C and D (CenA, CenB, CenC and CenD,
 15 respectively), exocellobiohydrolases A and B (CbhA and CbhB, respectively), and xylanases A and D (Cex and XylD, respectively) (see Wong *et al.* (1986) Gene, 44:315; Meinke *et al.* (1991) J. Bacteriol., 173:308; Coutinho *et al.*, (1991) Mol. Microbiol. 5:1221; Meinke *et al.*, (1993) Bacteriol., 175:1910; Meinke *et al.*, (1994) Mol. Microbiol.,
 20 12:413; Shen *et al.*, Biochem. J., in press; O'Neill *et al.*, (1986) Gene, 44:325; and Millward-Sadler *et al.*, (1994) Mol. Microbiol., 11:375). All are modular proteins of varying degrees of complexity, but with two features in common: a catalytic domain (CD) and a cellulose-binding domain (CBD) which can function independently (see Millward-Sadler *et al.*,
 25 *et al.*, (1994) Mol. Microbiol., 11:375; Gilkes *et al.*, (1988) J. Biol. Chem., 263:10401; Meinke *et al.*, (1991) J. Bacteriol., 173:7126; and Coutinho *et al.*, (1992) Mol. Microbiol., 6:1242). In four of the enzymes, CenB, CenD, CbhA and CbhB, fibronectin type III (Fn3) repeats separate the N-terminal CD from the C-terminal CBD. The CDs

of the enzymes come from six of the families of glycoside hydrolases (see Henrissat (1991) *Biochem. J.*, 280:309; and Henrissat *et al.*, (1993) *Biochem. J.*, 293:781); all of the enzymes have an N- or C-terminal CBD or CBDs (see Tomme *et al.*, *Adv. Microb. Physiol.*, in press); CenC has
 5 tandem CBDs from family IV at its N-terminus; CenB and XylD each have a second, internal CBD from families III and II, respectively. Cex and XylD are clearly xylanases; however, Cex, but not XylD, has low activity on cellulose. Nonetheless, like several other bacterial xylanases (see Gilbert *et al.*, (1993) *J. Gen. Microbiol.*, 139:187), they have
 10 CBDs. *C. fimi* probably produces other β -1,4-glycanases. Similar systems are produced by related bacteria (see Wilson (1992) *Crit. Rev. Biotechnol.*, 12:45; and Hazlewood *et al.*, (1992) *J. Appl. Bacteriol.*, 72:244). Unrelated bacteria also produce glycanases; *Clostridium thermocellum*, for example, produces twenty or more β -1,4-glycanases
 15 (see Beguin *et al.*, (1992) *FEMS Microbiol. Lett.*, 100:523). The CBD derived from *C. fimi* endoglucanase C N1, is the only protein known to bind soluble cellosaccharides and one of a small set of proteins that are known to bind any soluble polysaccharides.

Examples of suitable binding domains are shown in Figure 1 of
 20 PCT/CA97/00033 (WO 97/26358), which presents an alignment of binding domains from various enzymes that bind to polysaccharides and identifies amino acid residues that are conserved among most or all of the enzymes. This information can be used to derive a suitable oligonucleotide probe using methods known to those of skill in the art.
 25 The probes can be considerably shorter than the entire sequence but should at least be 10, preferably at least 14, nucleotides in length. Longer oligonucleotides are useful, up to the full length of the gene, preferably no more than 500, more preferably no more than 250, nucleotides in length. RNA or DNA probes can be used. In use, the probes are typically labeled

in a detectable manner, for example, with ^{32}P , ^3H , biotin, avidin or other detectable reagents, and are incubated with single-stranded DNA or RNA from the organism in which a gene is being sought. Hybridization is detected by means of the label after the unhybridized probe has been
5 separated from the hybridized probe. The hybridized probe is typically immobilized on a solid matrix such as nitrocellulose paper. Hybridization techniques suitable for use with oligonucleotides are well known to those skilled in the art. Although probes are normally used with a detectable label that allows easy identification, unlabeled oligonucleotides are also
10 useful, both as precursors of labeled probes and for use in methods that provide for direct detection of double-stranded DNA (or DNA/RNA). Accordingly, the term "oligonucleotide probe" refers to both labeled and unlabeled forms.

Generally, the binding domains identified by probing nucleic acids
15 from an organism of interest will show at least about 40 % identity (including as appropriate allowances for conservative substitutions, gaps for better alignment and the like) to the binding region or regions from which the probe was derived and will bind to a soluble β -1,4 glucan with a K_d of $\geq 10^3 \text{ M}^{-1}$. More preferably, the binding domains will be at least
20 about 60 % identical, and most preferably at least about 70 % identical to the binding region used to derive the probe. The percentage of identity will be greater among those amino acids that are conserved among polysaccharidase binding domains. Analyses of amino acid sequence comparisons can be performed using programs in PC/Gene
25 (IntelliGenetics, Inc.). PCLUSTAL can be used for multiple sequence alignment and generation of phylogenetic trees.

In order to isolate the polysaccharide binding protein or a polysaccharide binding domain from an enzyme or a cluster of enzymes that binds to a polysaccharide, several genetic approaches can be used.

One method uses restriction enzymes to remove a portion of the gene that codes for portions of the protein other than the binding portion thereof. The remaining gene fragments are fused with expression control sequences to obtain a mutated gene that encodes a truncated protein.

5 Another method involves the use of exonucleases such as *Bal31* to systematically delete nucleotides either externally from the 5' and the 3' ends of the DNA or internally from a restricted gap within the gene. These gene deletion methods result in a mutated gene encoding a shortened protein molecule which can then be evaluated for substrate or
10 polysaccharide binding ability.

Any cellulose binding protein or cellulose binding domain may be used in the present invention. The term "cellulose binding protein" ("CBP") refers to any protein or polypeptide which specifically binds to cellulose. The cellulose binding protein may or may not have cellulose or cellulolytic
15 activity. The term "cellulose binding domain" ("CBD") refers to any protein or polypeptide which is a region or portion of a larger protein, said region or portion binds specifically to cellulose. The cellulose binding domain (CBD) may be a part or portion of a cellulase, xylanase or other polysaccharidase, e.g., a chitinase, etc., a sugar binding protein such as
20 maltose binding protein, or scaffoldin such as CbpA of *Clostridium celluvorans*, etc. Many cellulases and hemicellulases (e.g. xylanases and mannases) have the ability to associate with cellulose. These enzymes typically have a catalytic domain containing the active site for substrate hydrolysis and a carbohydrate-binding domain or cellulose-binding domain
25 for binding cellulose. The CBD may also be from a non-catalytic polysaccharide binding protein. To date, more than one hundred cellulose-binding domains (CBDs) have been classified into at least thirteen families designated I-XIII (Tomme *et al.* (1995) "Cellulose Binding Domains: Classification and Properties", in ACS Symposium Series 618 Enzymatic

Degradation and Insoluble Carbohydrates, pp. 142-161, Saddler and Penner eds., American Chemical Society, Washington, D.C. (Tomme I) ; Tomme *et al.* Adv. Microb. Physiol. (1995) 37:1 (Tomme II); and Smant *et al.*, Proc. Natl. Acad. Sci U.S.A. (1998) 95:4906-4911, all of which are
5 incorporated herein by reference). Any of the CBDs described in Tomme I or II or any variants thereof, any other presently known CBDs or any new CBDs which may be identified can be used in the present invention. As an illustrative, but in no way limiting example, the CBP or CBD can be from a bacterial, fungal, slime mold, or nematode protein or polypeptide. For a
10 more particular illustrative example, the CBD is obtainable from *Clostridium cellulovorans*, *Clostridium cellulovorans*, or *Cellulomonas fimi* (e.g., CenA, CenB, CenD, Cex). In addition, the CBD may be selected from a phage display peptide or peptidomimetic library, random or otherwise, using e.g., cellulose as a screening agent. (See Smith Science
15 (1985) 228:1315-1317 and Lam, Nature (1991) 354:82-84). Furthermore, the CBD may be derived by mutation of a portion of a protein or polypeptide which binds to a polysaccharide other than cellulose (or hemicellulose) but also binds cellulose, such as a chitinase, which specifically binds chitin, or a sugar binding protein such as maltose binding
20 protein, rendering said portion capable of binding to cellulose. In any event, the CBD binds cellulose or hemicellulose. Shoseyov and Doi (Proc. Natl. Acad. Sci. USA (1990) 87:2192-2195) isolated a unique cellulose-binding protein (CbpA) from the cellulose "complex" of the cellulolytic bacterium *Clostridium cellulovorans*. This major subunit of the cellulose
25 complex was found to bind to cellulose, but had no hydrolytic activity, and was essential for the degradation of crystalline cellulose. The CbpA gene has been cloned and sequenced (Shoseyov *et al.* Proc. Natl. Acad. Sci. USA (1992) 89:3483-3487). Using PCR primers flanking the cellulose-binding domain of CbpA, the latter was successfully cloned into an

overexpression vector that enabled overproduction of the approximately 17 kDa CBD in *Escherichia coli*. The recombinant CBD exhibits very strong affinity to cellulose and chitin (U.S. Pat. No. 5,496,934; Goldstein *et al.*, J. Bacteriol. (1993) 175:5762; PCT International Publication WO 94/24158, all are incorporated by reference as if fully set forth herein).

In recent years, several CBDs have been isolated from different sources. Most of these have been isolated from proteins that have separate catalytic, i.e., cellulose and cellulose binding domains, and only two have been isolated from proteins that have no apparent hydrolytic activity but possess cellulose-binding activity (Goldstein *et al.* J. Bacteriol. (1993) 175:5762-5768; Morag *et al.* Appl. (1995) Environ. Microbiol. 61:1980-1986).

Recombinant proteins:

Any protein for which a gene is known or can be isolated can be used as the recombinant protein and be fused to the cellulose binding peptide according to the present invention. Advantageously, the recombinant protein is of a commercial value. A non-exhaustive list of recombinant proteins which can be manufactured utilizing the process of the present invention and their uses follows.

Thus, for example, glucoamylases and glucose isomerases are used in the food processing industry to convert starch to high fructose corn syrup.

Another useful class of enzymes are proteinases, which are used for the hydrolysis of high molecular weight proteins and which are further used in combination with detergents in cleaning applications, in leather manufacturing processes, in the food industry, and in the manufacture of alcoholic beverages.

Enzymes known as pectinesterases, and several related enzymes, are used for pectin hydrolysis in the food industry.

A class of enzymes known as lipases are used for the cleavage of ester linkages in triglycerides, and are used both in the food industry and for effluent treatment.

The enzyme beta-galactosidase is used industrially for the hydrolysis of whey lactose.

An enzyme known as thermolysin is used in the production of the artificial sweetener aspartame.

An enzyme known as sulphydryl oxidase is used in the reduction of the cooked flavor of milk.

Enzymes known as catalases are used to remove hydrogen peroxides from milk, cheese, and egg processing, and are further used in the sterilization and oxidation of plastics and rubbers.

Heparinases are useful for the production of heparin and heparan sulfate oligosaccharides.

Other proteins, in addition to enzymes, are those which have affinities to other compounds. For example, bacteria, fungi, plants and animals all contain a large number of proteins that exhibit specific interactions with agents such as metal ions and toxic compounds, and have high affinities for such agents.

A class of proteins known as metalloproteins contain prosthetic groups that bind specifically to metal ions. An example of such a prosthetic group is the porphyrin group in hemoglobin. Some other examples of metal ion binding proteins include parvalbumin, which binds to calcium, and metallothionin, an animal protein that binds large amounts of metal ions, especially zinc. Such metal absorptive proteins could also be used for purification in industrial processes.

It is also envisioned that streams of flowing material could be degraded by microbial enzymes. It is known that certain pollutants, whether natural or synthetic, and certain pesticides and other durable

organic compounds in the environment can be degraded (inactivated) or converted into useful compounds by microbial enzymes.

It is known, for example, that some microorganisms, for example *Pseudomonas putida*, possessed dehalogenases that are capable of
5 degrading certain pesticides and herbicides, and rendering them less toxic. Similarly, hydrolysis of organophosphate insecticides have been observed by microbial enzymes.

It is also possible to produce antibodies within plant cells. The antibodies can include monoclonal antibodies or fragments thereof having
10 at least a portion of an antigen binding region, including immunoactive entities such as Fv, F(abl)2, Fab fragments (Harlow and Lane, 1988 Antibody, Cold Spring Harbor), single chain antibodies (U.S. Pat. No. 4,946,778), chimeric or humanized antibodies (Morrison *et al.* Proc. Natl. Acad. Sci. USA (1984) 81:6851; Neuberger *et al.* Nature (1984) 312:604-
15 608) and complementarily determining regions (CDR).

Another class of proteins are those that bind to antibodies, such as protein-A, protein-G, protein-L and their mutants.

It is also possible to produce protein antibiotics or peptides such as lysozyme or therapeutic proteins which might assist in healing processes,
20 for example, certain wound healing peptides, growth factors and hormones. Proteins such as HSA can also be produced.

Another class of proteins include proteins such as agglutinin, zein, silk, elastine proteins as well as COMP, JUN, FOS and other proteins that may form stable protein-protein interactions such as coiled-coil interactions
25 that may be useful for production of protein fibers.

Another example is the production of animal feed enzymes. Phytase from *Aspergillus niger*, for example, increases the availability of phosphorus from feed for monogastric animals by releasing phosphate from the substrate phytic acid, therefore reducing the need for costly phosphorus

supplements. A phytase cDNA was constitutively expressed in transgenic tobacco (*Nicotiana tabacum*) plants (Verwoerd *et al.*, Plant. Physiol. (1995) 109:1199-205). Soybean plants transformed with a fungal phytase gene improve phosphorus availability whereas excretion was decreased for
 5 broilers. It appears that phytase can improve growth performance of broilers fed low phosphorous diets when provided either as a commercial supplement or in the form of transformed seeds (Denbow *et al.*, Poul. Sci. (1989) 77:878-881).

Other recombinant proteins of interest, will for the most part be
 10 mammalian proteins, and will include blood proteins, such as serum albumin, Factor VII, Factor VIIIc, Factor VIIIvW, Factor IX, Factor X, tissue plasminogen factor, Protein C, von Willebrand factor, antithrombin III, erythropoietin, colony stimulating factors, such as G-, M-, GM-, cytokines, such as interleukins 1-11, integrins, addressins, selectins, homing
 15 receptors, surface membrane proteins, such as surface membrane protein receptors, T cell receptor units, immunoglobulins (as further detailed above with respect to antibodies), soluble major histocompatibility complex antigens, structural proteins, such as collagen, fibrin, elastin, tubulin, actin, and myosin, growth factor receptors, growth factors, growth hormone, cell
 20 cycle proteins, vaccines, fibrinogen, thrombin, cytokines and hyaluronidase. Additional examples include chymosin, polymerases, saccharidases, dehydrogenases, nucleases, oxido reductases such as fungal peroxidases and lactases, xylanases, rennin, horse radish peroxidase, amylases and soil remediation enzymes.

25 The genes encoding all of the above listed proteins have been isolated and as such these proteins are readily available for recombinant expression and production according to the teachings of the present invention. It will be appreciated that new genes encoding an ever growing spectrum of proteins are continuously discovered and isolated, rendering

such genes available for molecular manipulation and recombinant expression. There is thus no intention to limit the recombinant protein produced utilizing the method of the present invention to any specific protein or list of proteins.

5 ***Cellulose binding peptide-recombinant protein fusions:***

The fusion of two proteins for which genes has been isolated is well known and practiced in the art. Such fusion involves the joining together of heterologous nucleic acid sequences, in frame, such that translation thereof results in the generation of a fused protein product or a fusion proteins.

10 Methods, such as the polymerase chain reaction (PCR), restriction, nuclease digestion, ligation, synthetic oligonucleotides synthesis and the like are typically employed in various combinations in the process of generating fusion gene constructs. One ordinarily skilled in the art can readily form such constructs for any pair or more of individual proteins. Interestingly, in
15 most cases where such fusion or chimera proteins are produced, and in all cases where one of the proteins was a cellulose binding peptide, both the former and the latter retained their catalytic activity or function.

For example, Greenwood *et al.* (1989, FEBS Lett. 224:127-131) fused the cellulose binding region of *Cellulomonas fimi* endoglucanase to
20 the enzyme alkaline phosphatase. The recombinant fusion protein retained both its phosphatase activity and the ability to bind to cellulose. For more descriptions of cellulose binding fusion proteins, see U.S. Patent No. 5,137,819 issued to Kilburn *et al.*, and U.S. Patent No. 5,719,044 issued to Shoseyov *et al.* both incorporated by reference herein. See also U.S. Pat.
25 No. 5,474,925. All of which are incorporated herein by reference.

The recombinant protein immobilized via its fused counterpart to the cellulosic matter can be released from the plant derived cellulosic matter by cleavage thereof, e.g., by proteolysis, using either a nonspecific general protease such as proteinase K or trypsin, or a specific protease as further

detailed hereinunder. For example, release can be effected by treatment with proteinase K at a concentration of about 50 µg/ml for about 20 minutes at about 37 °C (Din *et al.* Bio/Technology (1991) 9:1096-1099).

Inclusion of a dedicated cleavage site:

5 According to a preferred embodiment of the present invention the fusion protein includes the recombinant protein and the cellulose binding peptide separated therebetween via a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence..

10 As used herein in the specification and in the claims section that follows, the phrase "unique amino acid sequence recognizable and digestible by a protease" includes a protease recognition sequence which is both recognizable and readily accessible to a protease. Thus, the unique sequence can be a solitary sequence (i.e., which does not appear in the
15 recombinant protein and optionally also not in the cellulose binding peptide) or alternatively, the sole sequence of several similar sequences which is not sequestered from the protease due to the tertiary structure of the recombinant protein and optionally the cellulose binding peptide. In both these cases proteolysis will release the recombinant protein from the fusion
20 protein cellulosic matter complex.

As used herein in the specification and in the claims section that follows, the phrase "controllable intervening protein sequence" includes unique amino acid sequences capable of excision from or cleavage of a peptide under predetermined conditions in cis or in trans, e.g., increase in
25 temperature, exposure to light, unblocking of amino acid residues by dephosphorylation and treatment with chemical reagents or deglycosylation, examples include proteolysis effected under acidic conditions (HCl, e.g., to cleave between Asp and Pro) and proteolysis effected by a proteolyzing reagent, such as CNBr to cleave downstream of Met, all as known in the art

and/or as further described in U.S. Pat. No. 5,834,247, which is incorporated herein by reference.

Thus, according to an aspect of the present invention there is provided a composition of matter comprising (a) a plant derived cellulosic matter; and (b) a fusion protein including a recombinant protein and a cellulose binding peptide separated therebetween via a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence, wherein the fusion protein is complexed to the plant derived cellulosic matter by affinity binding via the cellulose binding peptide.

Nucleic acid molecules which can be used according to preferred embodiments of the present invention to express the fusion protein in plant cells would therefore include a heterologous nucleic acid sequence including (i) a first sequence encoding a cellulose binding peptide; (ii) a second sequence encoding a recombinant protein, wherein the first and second sequences are joined together in frame in either orientation; and (iii) a third sequence encoding a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence, the third sequence is between and in frame with the first and second sequences.

Thus, specific cleavage can be used to release the recombinant protein from the fusion protein cellulosic matter complex. For example, one can include a protease recognition site or a chemical cleavage site between the recombinant protein and the cellulose binding peptide. Examples of recognition sites include those for collagenase, thrombin, enterokinase, and Factor X_a which are cleaved specifically by the respective enzymes. Chemical cleavage sites sensitive, for example, to low pH or cyanogen bromide, can also be used.

Where cleavage is used, the recombinant protein can be cleaved readily from the cellulosic matter by the use of a protease specific for a sequence present therebetween and the cellulose binding peptide.

It will be appreciated in this respect that four main classes of specific proteases are known, including (i) cysteine proteases, including cathepsin B and L; (ii) aspartyl protease cathepsin D; (iii) serine proteases including plasmin, tissue-type plasminogen activator (tPA) and urokinase-type plasminogen activator (uPA), and (iv) matrix metalloproteinases (MMPs), including collagenases, gelatinases A and B (MMP2 and MMP9) and stromelysin (MMP3). Members of these protease families are commercially available and their recognition sequences known. As such, these proteases can be used to implement the step of releasing the recombinant protein from the plant derived cellulosic matter while implementing the process according to the present invention.

Genetically modified plant material:

According to an aspect of the present invention there is provided a nucleic acid molecule comprising (a) a promoter sequence for directing protein expression in plant cells; and (b) a heterologous nucleic acid sequence as further detailed herein, wherein, the heterologous nucleic acid sequence is down stream the promoter sequence, such that expression of the heterologous nucleic acid sequence is effectable by the promoter sequence. Such a nucleic acid molecule needs to be effectively introduced into plant cells, so as to genetically modify the plant.

There are various methods of introducing foreign genes into both monocotyledonous and dicotyledonous plants (Potrykus, I., *Annu. Rev. Plant. Physiol., Plant. Mol. Biol.* (1991) 42:205-225; Shimamoto *et al.*, *Nature* (1989) 338:274-276). The principle methods of causing stable integration of exogenous DNA into plant genomic DNA include two main approaches:

(i) *Agrobacterium*-mediated gene transfer: Klee *et al.* (1987) Annu. Rev. Plant Physiol. 38:467-486; Klee and Rogers in Cell Culture and Somatic Cell Genetics of Plants, Vol. 6, Molecular Biology of Plant Nuclear Genes, eds. Schell, J., and Vasil, L. K., Academic Publishers, San Diego, Calif. (1989) p. 2-25; Gatenby, in Plant Biotechnology, eds. Kung, S. and Arntzen, C. J., Butterworth Publishers, Boston, Mass. (1989) p. 93-112.

(ii) direct DNA uptake: Paszkowski *et al.*, in Cell Culture and Somatic Cell Genetics of Plants, Vol. 6, Molecular Biology of Plant Nuclear Genes eds. Schell, J., and Vasil, L. K., Academic Publishers, San Diego, Calif. (1989) p. 52-68; including methods for direct uptake of DNA into protoplasts, Toriyama, K. *et al.* (1988) Bio/Technology 6:1072-1074. DNA uptake induced by brief electric shock of plant cells: Zhang *et al.* Plant Cell Rep. (1988) 7:379-384. Fromm *et al.* Nature (1986) 319:791-793. DNA injection into plant cells or tissues by particle bombardment, Klein *et al.* Bio/Technology (1988) 6:559-563; McCabe *et al.* Bio/Technology (1988) 6:923-926; Sanford, Physiol. Plant. (1990) 79:206-209; by the use of micropipette systems: Neuhaus *et al.*, Theor. Appl. Genet. (1987) 75:30-36; Neuhaus and Spangenberg, Physiol. Plant. (1990) 79:213-217; or by the direct incubation of DNA with germinating pollen, DeWet *et al.* in Experimental Manipulation of Ovule Tissue, eds. Chapman, G. P. and Mantell, S. H. and Daniels, W. Longman, London, (1985) p. 197-209; and Ohta, Proc. Natl. Acad. Sci. USA (1986) 83:715-719.

25 The *Agrobacterium* system includes the use of plasmid vectors that contain defined DNA segments that integrate into the plant genomic DNA. Methods of inoculation of the plant tissue vary depending upon the plant species and the *Agrobacterium* delivery system. A widely used approach is the leaf disc procedure which can be performed with any tissue explant that

provides a good source for initiation of whole plant differentiation. Horsch *et al.* in Plant Molecular Biology Manual A5, Kluwer Academic Publishers, Dordrecht (1988) p. 1-9. The *Agrobacterium* system is especially viable in the creation of transgenic dicotyledenous plants.

5 There are various methods of direct DNA transfer into plant cells. In electroporation, the protoplasts are briefly exposed to a strong electric field. In microinjection, the DNA is mechanically injected directly into the cells using very small micropipettes. In microparticle bombardment, the DNA is adsorbed on microprojectiles such as magnesium sulfate crystals or tungsten
10 particles, and the microprojectiles are physically accelerated into cells or plant tissues.

Following transformation plant propagation is exercised. The most common method of plant propagation is by seed. Regeneration by seed propagation, however, has the deficiency that due to heterozygosity there is
15 a lack of uniformity in the crop, since seeds are produced by plants according to the genetic variances governed by Mendelian rules. Basically, each seed is genetically different and each will grow with its own specific traits. Therefore, it is preferred that the transgenic plant be produced such that the regenerated plant has the identical traits and characteristics of the
20 parent transgenic plant, e.g., a reproduction of the fusion protein. Therefore, it is preferred that the transgenic plant be regenerated by micropropagation which provides a rapid, consistent reproduction of the transgenic plants.

Micropropagation is a process of growing new generation plants
25 from a single piece of tissue that has been excised from a selected parent plant or cultivar. This process permits the mass reproduction of plants having the preferred tissue expressing the fusion protein. The new generation plants which are produced are genetically identical to, and have all of the characteristics of, the original plant. Micropropagation allows

mass production of quality plant material in a short period of time and offers a rapid multiplication of selected cultivars in the preservation of the characteristics of the original transgenic or transformed plant. The advantages of cloning plants are the speed of plant multiplication and the quality and uniformity of plants produced.

Micropropagation is a multi-stage procedure that requires alteration of culture medium or growth conditions between stages. Thus, the micropropagation process involves four basic stages: Stage one, initial tissue culturing; stage two, tissue culture multiplication; stage three, differentiation and plant formation; and stage four, greenhouse culturing and hardening. During stage one, initial tissue culturing, the tissue culture is established and certified contaminant-free. During stage two, the initial tissue culture is multiplied until a sufficient number of tissue samples are produced to meet production goals. During stage three, the tissue samples grown in stage two are divided and grown into individual plantlets. At stage four, the transgenic plantlets are transferred to a greenhouse for hardening where the plants' tolerance to light is gradually increased so that it can be grown in the natural environment.

The basic bacterial/plant vector construct will preferably provide a broad host range prokaryote replication origin; a prokaryote selectable marker; and, for *Agrobacterium* transformations, T DNA sequences for *Agrobacterium*-mediated transfer to plant chromosomes. Where the heterologous sequence is not readily amenable to detection, the construct will preferably also have a selectable marker gene suitable for determining if a plant cell has been transformed. A general review of suitable markers for the members of the grass family is found in Wilmink and Dons, Plant Mol. Biol. Repr. (1993) 11:165-185.

Sequences suitable for permitting integration of the heterologous sequence into the plant genome are also recommended. These might

include transposon sequences and the like for homologous recombination as well as Ti sequences which permit random insertion of a heterologous expression cassette into a plant genome.

Suitable prokaryote selectable markers include resistance toward
 5 antibiotics such as ampicillin or tetracycline. Other DNA sequences encoding additional functions may also be present in the vector, as is known in the art.

The constructs of the subject invention will include an expression
 cassette for expression of the fusion protein of interest. Usually, there will
 10 be only one expression cassette, although two or more are feasible. The recombinant expression cassette will contain in addition to the heterologous sequence one or more of the following sequence elements, a promoter region, plant 5' untranslated sequences, initiation codon depending upon whether or not the structural gene comes equipped with one, and a
 15 transcription and translation termination sequence. Unique restriction enzyme sites at the 5' and 3' ends of the cassette allow for easy insertion into a pre-existing vector.

Viral infected plant material:

Viruses are a unique class of infectious agents whose distinctive
 20 features are their simple organization and their mechanism of replication. In fact, a complete viral particle, or virion, may be regarded mainly as a block of genetic material (either DNA or RNA) capable of autonomous replication, surrounded by a protein coat and sometimes by an additional membranous envelope such as in the case of alpha viruses. The coat
 25 protects the virus from the environment and serves as a vehicle for transmission from one host cell to another.

Viruses that have been shown to be useful for the transformation of plant hosts include CaV, TMV and BV. Transformation of plants using plant viruses is described in U.S. Pat. No. 4,855,237 (BGV), EP-A 67,553

(TMV), Japanese Published Application No. 63-14693 (TMV), EPA 194,809 (BV), EPA 278,667 (BV); and Gluzman, Y. *et al.*, Communications in Molecular Biology: Viral Vectors, Cold Spring Harbor Laboratory, New York, pp. 172-189 (1988). Pseudovirus particles for use
5 in expressing foreign DNA in many hosts, including plants, is described in WO 87/06261.

Construction of plant RNA viruses for the introduction and expression of non-viral foreign genes in plants is demonstrated by the above references as well as by Dawson, W. O. *et al.*, Virology (1989) 172:285-
10 292; Takamatsu *et al.* EMBO J. (1987) 6:307-311; French *et al.* Science (1986) 231:1294-1297; and Takamatsu *et al.* FEBS Letters (1990) 269:73-76.

When the virus is a DNA virus, the constructions can be made to the virus itself. Alternatively, the virus can first be cloned into a bacterial
15 plasmid for ease of constructing the desired viral vector with the foreign DNA. The virus can then be excised from the plasmid. If the virus is a DNA virus, a bacterial origin of replication can be attached to the viral DNA, which is then replicated by the bacteria. Transcription and translation of this DNA will produce the coat protein which will encapsidate
20 the viral DNA. If the virus is an RNA virus, the virus is generally cloned as a cDNA and inserted into a plasmid. The plasmid is then used to make all of the constructions. The RNA virus is then produced by transcribing the viral sequence of the plasmid and translation of the viral genes to produce the coat protein(s) which encapsidate the viral RNA.

25 Construction of plant RNA viruses for the introduction and expression of non-viral foreign genes in plants is demonstrated by the above references as well as in U.S. Pat. No. 5,316,931

In one embodiment, a plant viral nucleic acid is provided in which the native coat protein coding sequence has been deleted from a viral

nucleic acid, a non-native plant viral coat protein coding sequence and a non-native promoter, preferably the subgenomic promoter of the non-native coat protein coding sequence, capable of expression in the plant host, packaging of the recombinant plant viral nucleic acid, and ensuring a systemic infection of the host by the recombinant plant viral nucleic acid, has been inserted. Alternatively, the coat protein gene may be inactivated by insertion of the non-native nucleic acid sequence within it, such that a fusion protein is produced. The recombinant plant viral nucleic acid may contain one or more additional non-native subgenomic promoters. Each non-native subgenomic promoter is capable of transcribing or expressing adjacent genes or nucleic acid sequences in the plant host and incapable of recombination with each other and with native subgenomic promoters. Non-native (foreign) nucleic acid sequences may be inserted adjacent the native plant viral subgenomic promoter or the native and a non-native plant viral subgenomic promoters if more than one nucleic acid sequence is included. The non-native nucleic acid sequences are transcribed or expressed in the host plant under control of the subgenomic promoter to produce the desired products.

In a second embodiment, a recombinant plant viral nucleic acid is provided as in the first embodiment except that the native coat protein coding sequence is placed adjacent one of the non-native coat protein subgenomic promoters instead of a non-native coat protein coding sequence.

In a third embodiment, a recombinant plant viral nucleic acid is provided in which the native coat protein gene is adjacent its subgenomic promoter and one or more non-native subgenomic promoters have been inserted into the viral nucleic acid. The inserted non-native subgenomic promoters are capable of transcribing or expressing adjacent genes in a plant host and are incapable of recombination with each other and with

native subgenomic promoters. Non-native nucleic acid sequences may be inserted adjacent the non-native subgenomic plant viral promoters such that said sequences are transcribed or expressed in the host plant under control of the subgenomic promoters to produce the desired product.

5 In a fourth embodiment, a recombinant plant viral nucleic acid is provided as in the third embodiment except that the native coat protein coding sequence is replaced by a non-native coat protein coding sequence.

The viral vectors are encapsidated by the coat proteins encoded by the recombinant plant viral nucleic acid to produce a recombinant plant virus. The recombinant plant viral nucleic acid or recombinant plant virus is used to infect appropriate host plants. The recombinant plant viral nucleic acid is capable of replication in the host, systemic spread in the host, and transcription or expression of foreign gene(s) in the host to produce the desired fusion protein.

15 ***Fusion protein compartmentalization - signal peptides:***

As already mentioned hereinabove, compartmentalization of the fusion protein is an important feature of the present invention because it allows undisturbed plant growth. Thus, according to one aspect of the present invention, the fusion protein is compartmentalized within cells of the plant or cultured plant cells, so as to be sequestered from cell walls of the cells of the plant or cultured plant cells.

The fusion protein can be compartmentalized within a cellular compartment, such as, for example, the cytoplasm, endoplasmic reticulum, golgi apparatus, oil bodies, starch bodies, chloroplastids, chloroplasts, 25 chromoplastids, chromoplasts, vacuole, lysosomes, mitochondria or the nucleus.

Accordingly, the heterologous sequence used while implementing the process according to this aspect of the present invention includes (i) a first sequence encoding a cellulose binding peptide; (ii) a second sequence

encoding a recombinant protein, wherein the first and second sequences are joined together in frame; and (iii) a third sequence encoding a signal peptide for directing a protein to a cellular compartment, the third sequence being upstream and in frame with the first and second sequences.

5 The following provides description of signal peptides which can be used to direct the fusion protein according to the present invention to specific cell compartments.

10 It is well-known that signal peptides serve the function of translocation of produced protein across the endoplasmic reticulum membrane. Similarly, transmembrane segments halt translocation and provide anchoring of the protein to the plasma membrane, see, Johnson *et al.* The Plant Cell (1990) 2:525-532; Sauer *et al.* EMBO J. (1990) 9:3045-3050; Mueckler *et al.* Science (1985) 229:941-945. Mitochondrial, nuclear, chloroplast, or vacuolar signals target expressed protein correctly into the
15 corresponding organelle through the secretory pathway, see, Von Heijne, Eur. J. Biochem. (1983) 133:17-21; Von Heijne, J. Mol. Biol. (1986) 189:239-242; Iturriaga *et al.* The Plant Cell (1989) 1:381-390; McKnight *et al.*, Nucl. Acid Res. (1990) 18:4939-4943; Matsuoka and Nakamura, Proc. Natl. Acad. Sci. USA (1991) 88:834-838. A recent book by Cunningham
20 and Porter (Recombinant proteins from plants, Eds. C. Cunningham and A.J.R. Porter, 1998 Humana Press Totowa, N.J.) describe methods for the production of recombinant proteins in plants and methods for targeting the proteins to different compartments in the plant cell. In particular, two chapters therein (14 and 15) describe different methods to introduce
25 targeting sequences that results in accumulation of recombinant proteins in compartments such as ER, vacuole, plastid, nucleus and cytoplasm. The book by Cunningham and Porter is incorporated herein by reference. Presently, the preferred site of accumulation of the fusion protein according to the present invention is the ER using signal peptide such as Cel 1 or the

rice amylase signal peptide at the N-terminus and an ER retaining peptide (HDEL, SEQ ID NO:1; or KDEL, SEQ ID NO:2) at the C-terminus.

Promoters and control of expression:

Any promoter which can direct the expression of the fusion protein according to the present invention can be utilized to implement the process of the instant invention, both constitutive and tissue specific promoters. According to presently preferred embodiment the promoter selected is constitutive, because such a promoter can direct the expression of higher levels of the fusion protein. In this respect the present invention offers a major advantage over the teachings of U.S. Pat. No. 5,474,925 in which only tissue specific and weak promoters can be employed because of the deleterious effect of the fusion protein described therein on cell wall development. The reason for which the present invention can utilize strong and constitutive promoters relies in the compartmentalization and sequestering approach which prohibits contact between the expressed fusion protein and the plant cell walls which such walls are developing.

Constitutive and tissue specific promoters, CaMV35S promoter (Odell *et al.* Nature (1985) 313:810-812) and ubiquitin promoter (Christensen and Quail, Transgenic research (1996) 5:213-218) are the most commonly used constitutive promoters in plant transformations and are the preferred promoters of choice while implementing the present invention.

In corn, within the kernel, proteins under the ubiquitin promoters, are preferentially accumulated in the germ (Kusnadi *et al.*, Biotechnol. Bioeng. (1998) 60:44-52). The amylose-extender (Ae) gene encoding starch-branching enzyme IIb (SBEIIb) in maize is predominantly expressed in endosperm and embryos during kernel development (Kim *et al.* Plant. Mol. Biol. (1998) 38:945-956). A starch branching enzyme (SBE) showed promoter activity after it was introduced into maize endosperm suspension cells by particle bombardment (Kim *et al.* Gene (1998) 216:233-243). In

transgenic wheat it has been shown that a native HMW-GS gene promoter can be used to obtain high levels of expression of seed storage and, potentially, other proteins in the endosperm (Blechl and Anderson, Nat. Biotechnol. (1996) 14:875-9). Polygalacturonase (PG) promoter was shown

5 to confer high levels of ripening-specific gene expression in tomato (Nicholass *et al.* Plant. Mol. Biol. (1995) 28:423-435). The ACC oxidase promoter (Blume and Grierson, Plant. J. (1997) 12:731-746) represents a promoter from the ethylene pathway and shows increased expression during fruit ripening and senescence in tomato. The promoter for tomato 3-

10 hydroxy-3-methylglutaryl coenzyme A reductase gene accumulates to high level during fruit ripening (Daraselia *et al.* Plant. Physiol. (1996) 112:727-733). Specific protein expression in potato tubers can be mediated by the patatin promoter (Sweetlove *et al.* Biochem. J. (1996) 320:487-492). Protein linked to a chloroplast transit peptide changed the protein content in

15 transgenic soybean and canola seeds when expressed from a seed-specific promoter (Falco *et al.* Biotechnology (NY) (1995) 13:577-82). The seed specific bean phaseolin and soybean beta-conglycinin promoters are also suitable for the latter example (Keeler *et al.* Plant. Mol. Biol. (1997) 34:15-29). Promoters that are expressed in plastids are also suitable in

20 conjunction with plastid transformation.

Each of these promoters can be used to implement the process according to the present invention.

Thus, the plant promoter employed can a constitutive promoter, a tissue specific promoter, an inducible promoter or a chimeric promoter.

25 Examples of constitutive plant promoters include, without being limited to, CaMV35S and CaMV19S promoters, FMV34S promoter, sugarcane bacilliform badnavirus promoter, CsVMV promoter, *Arabidopsis* ACT2/ACT8 actin promoter, *Arabidopsis* ubiquitin UBQ1 promoter, barley leaf thionin BTH6 promoter, and rice actin promoter.

Examples of tissue specific promoters include, without being limited to, bean phaseolin storage protein promoter, DLEC promoter, PHS β promoter, zein storage protein promoter, conglutin gamma promoter from soybean, AT2S1 gene promoter, ACT11 actin promoter from *Arabidopsis*,
 5 napA promoter from *Brassica napus* and potato patatin gene promoter.

The inducible promoter is a promoter induced by a specific stimuli such as stress conditions comprising, for example, light, temperature, chemicals, drought, high salinity, osmotic shock, oxidant conditions or in case of pathogenicity and include, without being limited to, the light-
 10 inducible promoter derived from the pea rbcS gene, the promoter from the alfalfa rbcS gene, the promoters DRE, MYC and MYB active in drought; the promoters INT, INPS, prxEa, Ha hsp17.7G4 and RD21 active in high salinity and osmotic stress, and the promoters hsr303J and str246C active in pathogenic stress.

15 ***Expression follow up:***

Expression of the fusion protein can be monitored by a variety of methods. For example, ELISA or western blot analysis using antibodies specifically recognizing the recombinant protein or its cellulose binding peptide counterpart can be employed to qualitatively and/or quantitatively
 20 monitor the expression of the fusion protein in the plant. Alternatively, the fusion protein can be monitored by SDS-PAGE analysis using different staining techniques, such as, but not limited to, coomassie blue or silver staining. Other methods can be used to monitor the expression level of the RNA encoding for the fusion protein. Such methods include RNA
 25 hybridization methods, e.g., Northern blots and RNA dot blots.

Binding of the fusion protein to the plant derived cellulosic matter:

When sufficient expression has been detected, binding of the fusion protein to the plant derived cellulosic matter is effected. Such binding can be achieved, for example, as follows. Whole plants, plant derived tissue or

cultured plant cells are homogenized by mechanical method in the presence or absence of a buffer, such as, but not limited to, PBS. The fusion protein is therefore given the opportunity to bind to the plant derived cellulosic matter. Buffers that may include salts and/or detergents at optimal concentrations may be used to wash non specific proteins from the cellulosic matter.

Extraction and purification:

In general, a recent book by Cunningham and Porter (Recombinant proteins from plants, Eds. C. Cunningham and A.J.R. Porter, 1998 Humana Press Totowa, N.J.) describes methods for the production of recombinant proteins in plants including methods for extraction of the proteins from the plants. The methods used herein for extraction of proteins from plants are similar, however the ability of the fusion protein to bind to cellulose dictates its fate, unless extraction is done under condition in which the cellulose binding peptide do not bind to cellulose, for example, pH higher than 10 (for most CBDs) or high concentration of glucose or cellobiose (200 mM or higher) for family IX CBDs. If the initial extraction is conducted under conditions that prevent binding, the supernatant is cleared from the cellulosic matter and then the solution is brought by either dilution, dialysis or pH correction, if necessary, to a condition that enables binding, after which cellulose is added in a batch or the solution is loaded on a cellulose column. Cellulose affinity purification is conducted as described, for example, in U.S. Pat. Nos. 5,719,044; 5,670,623; 5,856,021; 5,137,819; 5,202,247; 5,340,731; and 5,474,925; and U.S. Pat. applications Nos. 08/788,621; and 08/788,622; EP 0 381 719 B1, and EP application No. 93907724.4. Alternatively, the extraction solution provides conditions that favor binding to the plant derived cellulosic matter.

In any case, while the fusion protein is bound to cellulose, further whases can be employed for further removal of unbound proteins,

conditions which dissociate such binding or proteolytic cleavage can be used to isolate the fusion protein itself, or proteolytic cleavage can be used to isolate the recombinant protein, all as further detailed hereinabove.

5 Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which is not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims
10 section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together
15 with the above descriptions, illustrate the invention in a non limiting fashion. Generally, the nomenclature used herein and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification.
20 Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturers' specifications. These techniques and various other techniques are generally performed according to Sambrook *et al.*, molecular Cloning - A Laboratory Manual, Cold Spring Harbor Laboratory, Cold
25 Spring Harbor, N.Y. (1989). Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein, as well as that contained in the Manual by Sambrook *et al.*, is incorporated herein by reference.

EXAMPLE 1

Expression of CBD-Protein-L in transgenic potato tubers

Constructs:

5 A vector containing the class-I patatin gene B33 promoter (Olesinski *et al.*, 1996, Plant Physiol. 111:541-550), fused to the cell signal sequence (Shani *et al.*, 1997, Plant Mol. Biol. 34:837-842), protein-L (hereinafter ProtL, Nilson *et al.*, 1993, J. Immunol. Meth. 164:33-40), CBD cex sequence (Gilkes *et al.*, 1991, Microbiol. Rev. 55:303-315) and HDEL
10 (SEQ ID NO:1) retaining peptide sequence was constructed as follows.

A DNA fragment encoding the cell signal peptide (nucleotide 1-105 of SEQ ID NO:3) was generated by PCR using the following primers: No. 1 (BglII) 5'-AAAAAGATCTATGGCGCGAAAATCACTAA-3' (SEQ ID NO:4) and No. 2 (XbaI) 5'-AAAATCTAGATTTACGGAGAGCGTCGCG-
15 3' (SEQ ID NO:5). A DNA fragment encoding the ProtL-CBDcex (nucleotides 3-1280 of SEQ ID NO:6) was generated by PCR using the following primers: No. 3 (XbaI) 5'-AAAATCTAGAATGGCG GCGGTAGAAAATAAAG-3' (SEQ ID NO:7); and No. 4 (HDEL, Stop and SalI) 5'-AAAAGTCGACTTAAAGTTCATCATGCTCGACGCCGA
20 CCGTGCA-3' (SEQ ID NO:8). The two fragment were digested with BglII, XbaI and SalI and ligated in one step into pUC19 (New England Biolabs, Beverly, Massachusetts) that was pre digested with BamHI and SalI. The primer for the c-terminal end of the ProtL-CBD contained the retaining peptide sequence HDEL (SEQ ID NO:1) and a stop codon in
25 frame. The DNA containing the Cell Signal-ProtL-CBD-HDEL (hereinafter, ProtL-CBD) fusion was excised using SmaI and SalI and was subcloned into the SmaI and SalI sites of the binary vector Bin19 (Bevan, 1984, Nuc. Acid Res. 12:8711-8721) under the class-I patatin gene B33 promoter (Olesinski *et al.*, 1996, Plant Physiol. 111:541-550).

Construction of transgenic plants:

The binary vector was mobilized into disarmed LB 4404 *Agrobacterium tumefaciens* (An, 1987, Meth. Enzymol. 153:292-305). Tuber discs transformation was performed with *Solanum tuberosum* cv Desiree plants as described previously (Olesinski *et al.*, 1996, 1996, Plant Physiol. 111:541-550). Regenerated transgenic plants were selected on kanamycin.

Analysis of transgenic plants:

Transformant plants were selected by rooting on kanamycin (100 mg/L). Growth conditions in growth room and green house were as described previously (Olesinski *et al.*, 1996, 1996, Plant Physiol. 111:541-550).

Transgenic plants transformed with the ProtL-CBD construct can be tested for expression of ProtL-CBD. Thus, for example, a Northern blot analysis of ProtL-CBD RNA expression can be carried out using the DNA fragment of SEQ ID NO:3 or a portion thereof as a probe. Furthermore, RT-PCR can be performed on RNA samples obtained from such transgenic plants using PCR primers No. 3 and No. 4 to amplify and detect the ProtL-CBD RNA. Additional methods for analysis of expression in transgenic plants include RNase protection assays. Expression of the ProtL-CBD fusion protein can alternatively or additionally be evaluated by monitoring the fusion protein itself using, for example, ELISA or Western blot protocols and anti-CBD or ProtL antibodies. Yet in addition, the ProtL-CBD fusion protein can be visualized by SDS-PAGE analysis using different staining techniques, such as coomassie blue or silver staining, immuno precipitation, enzyme linked immunoassays or CBD and ProtL binding assays. In addition, techniques such as *in situ* hybridization and immunostaining also may be used to detect the presence or expression of the

recombinant construct in specific plant organs and tissues. The methods for performing all of these assays are well known to those skilled in the art.

Although the invention has been described in conjunction with
5 specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims. All publications cited herein are incorporated by reference in their
10 entirety.

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Ziv Shani et al.
 - (ii) TITLE OF INVENTION: PROCESS OF EXPRESSING AND ISOLATING RECOMBINANT PROTEINS AND RECOMBINANT PROTEIN PRODUCTS FROM PLANTS, PLANT DERIVED TISSUES OR CULTURED PLANT CELLS
 - (iii) NUMBER OF SEQUENCES: 43
 - (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Mark M. Friedman c/o Anthony Castorina
 - (B) STREET: 2001 Jefferson Davis Highway, Suite 207
 - (C) CITY: Arlington
 - (D) STATE: Virginia
 - (E) COUNTRY: United States of America
 - (F) ZIP: 22202
 - (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: 1.44 megabyte, 3.5" microdisk
 - (B) COMPUTER: Twinhead@ Slimnote-890TX
 - (C) OPERATING SYSTEM: MS DOS version 6.2, Windows version 3.11
 - (D) SOFTWARE: Word for Windows version 2.0 converted to an ASCII file
 - (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
 - (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Friedman, Mark M.
 - (B) REGISTRATION NUMBER: 33,883
 - (C) REFERENCE/DOCKET NUMBER: 1079/5
 - (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 972-3-5625553
 - (B) TELEFAX: 972-3-5625554
 - (C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:
His Asp Glu Leu
4
- (2) INFORMATION FOR SEQ ID NO:2:
- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 4
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:
Lys Asp Glu Leu
4

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 150
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

ATGGCGCGAA AATCCCTAAT TTCCCGGTG ATTTGCTCG CCGTCTCTCT 50
 CTTCTCTCCG CCGATTACT CCGCCGGTCA CGATTACCGC GACGCTCTCC 100
 GTAAAAGCAT TCTCTCTTC GAAGGTCAAC GTTCCGGTAA ACTCCCTCCA 150

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AAAAAGATCT ATGGCGCGAA AATCACTAA 29

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AAAATCTAGA TTACGGAGA GCGTCGCG 29

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1305
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCATGGCGGC GGTAGAAAAT AAAGAAGAAA CACCAGAAAC ACCAGAACT 50
 GATTACAGAAG AAGAAGTAAC AATCAAAGCT AACCTAATCT TTGCAAAATGG 100
 AAGCACACAA ACTGCAGAAT TCAAAGGAAC ATTTGAAAAA GCAACATCAG 150
 AAGCTTATGA GTATGCAGAT ACITTGAAGA AAGACAATGG AGAATATACT 200
 GTAGATGTTG CAGATAAAGG TTATACTTTA AATATTAAAT TTGCTGGAAA 250
 AGAAAAACA CAGAAGAAC CAAAAGAAGA AGTTACTATT AAAGCAAACT 300
 TAATCTATGC AGATGGAAAA ACACAAACAG CAGAATTCAG AGGAACATTT 350
 GAAGAAGCAA CAGCAGAAGC ATACAGATAT GCAGATGCAT TAAAGAAGGA 400
 CAATGGAGAA TATACAGTAG ACGTTGCAGA TAAAGGTTAT ACTTTAAATA 450
 TTAAATTTGC TGGAAAAGAA AAAACACCAG AAGAACCAAA AGAAGAAGTT 500
 ACTATTAAAG CAAACTTAAT CTATGCAGAT GGAAAAACAC AAACAGCAGA 550
 ATTCAAAGGA ACATTTGAAG AAGCAACAGC AGAAGCATAC AGATATGCTG 600
 ACTTATTAGC AGCAAAAGAA AATGGTAAAT ATACAGTAGA CGTTGCAGAT 650
 AAAGGTTATA CTTTAAATAT TAAATTGCT GGAAAAGAAA AAACACCAGA 700
 AGAACCAAAA GAAGAAGTTA CTATTAAAGC AAACCTAATC TATGCAGATG 750
 GAAAACTCA AACAGCAGAG TTCAAAGGAA CATTTGCAGA AGCAACAGCA 800
 GAAGCATACA GATACGCTGA CTTATTAGCA AAAGAAAATG GTAAATATAC 850
 AGCAGACTTA GAAGATGTTG GATACACTAT TAATATTAGA TTTGCAGGTA 900
 AGAAAGTTGA CGAAAAACCA GAAGGGATCC CTCCGACGCC GACCCCGACT 950
 AGTGGTCCGG CCGGGTGCCA GGTGCTGTGG GCGTCAACC AGTGAACAC 1000
 CGGCTTCACC GCGAAOGTCA CCGTGAAGAA CACGTCTCTC GCTCCGGTAG 1050
 ACGGCTGGAC GCTCACGTTT AGCTTCCCGT CCGGCCAGCA GGTCAACCAG 1100

GCGTGGAGCT CGACGGTCAC GCAGTCCGGC TCGGCCGTGA CGGTCCGCAA 1150
 CGCCCCGTGG AACGGCTCGA TCCCGGCGGG CGGCACCGCG CAGTTCGGCT 1200
 TCAACGGCTC GCACACGGGC ACCAACGCGG CGCCGACGGC GTTCTCGCTC 1250
 AACGGCACGC CTTGCACGGT CGGCGTCGAG CACCACCACC ACCACCACCA 1300
 CCACT 1305

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AAAATCTAGA ATGGCGGCGG TAGAAAATAA AG 32

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

AAAAGTCGAC TTAAAGTTCA TCATGCTCGA CGCCGACCGT GCA 43

WHAT IS CLAIMED IS:

1. A process of expressing a recombinant protein in a plant and of isolating the recombinant protein from the plant, the process comprising the steps of:

- (a) providing a plant, a plant derived tissue or cultured plant cells expressing a fusion protein including the recombinant protein and a cellulose binding peptide being fused thereto, said fusion protein being compartmentalized within cells of said plant, plant derived tissue or cultured plant cells, so as to be sequestered from cell walls of said cells of said plant, plant derived tissue or cultured plant cells;
- (b) homogenizing said plant, plant derived tissue or cultured plant cells, so as to bring into contact said fusion protein with a plant derived cellulosic matter of said plant, plant derived tissue or cultured plant cells, to thereby effect affinity binding of said fusion protein via said cellulose binding peptide to said cellulosic matter, thereby obtaining a fusion protein cellulosic matter complex; and
- (c) isolating said fusion protein cellulosic matter complex.

2. The process of claim 1, further comprising the step of:

- (d) washing said fusion protein cellulosic matter complex, thereby removing endogenous plant proteins and other plant material therefrom.

3. The process of claim 2, further comprising the step of:

- (e) collecting said fusion protein cellulosic matter complex as a final product of the process.

4. The process of claim 2, further comprising the step of:

- (e) exposing said fusion protein cellulosic matter complex to conditions effective in dissociating said fusion protein from said cellulosic matter; and
- (f) isolating said fusion protein, thereby obtaining an isolated fusion protein.

5. The process of claim 4, wherein said conditions effective in dissociating said fusion protein from said cellulosic matter are selected from the group consisting of basic conditions, denaturative conditions and affinity displacement conditions.

6. The process of claim 4, further comprising the step of:

- (g) exposing said isolated fusion protein to conditions effective in digesting said fusion protein so as to release said recombinant protein therefrom, thereby obtaining a released recombinant protein.

7. The process of claim 6, wherein said conditions effective in digesting said fusion protein so as to release said recombinant protein therefrom are selected from the group consisting of proteolysis effected via a protease and proteolysis effected under predetermined cis or trans conditions for digesting a controllable intervening protein sequence.

8. The process of claim 4, further comprising the step of:

- (h) isolating said released recombinant protein.

9. The process of claim 2, further comprising the step of:
- (e) exposing said fusion protein cellulosic matter complex to conditions effective in digesting said fusion protein so as to release said recombinant protein therefrom, thereby obtaining a released recombinant protein.

10. The process of claim 6, wherein said conditions effective in digesting said fusion protein so as to release said recombinant protein therefrom are selected from the group consisting of proteolysis effected via a protease and proteolysis effected under predetermined cis or trans conditions for digesting a controllable intervening protein sequence.

11. The process of claim 4, further comprising the step of:
- (f) isolating said released recombinant protein.

12. A genetically modified or viral infected plant or cultured plant cells expressing a fusion protein including a recombinant protein and a cellulose binding peptide, said fusion protein being compartmentalized within cells of said plant or cultured plant cells, so as to be sequestered from cell walls of said cells of said plant or cultured plant cells.

13. The genetically modified or viral infected plant or cultured plant cells of claim 12, wherein expression of said fusion protein is under a control of a constitutive or tissue specific plant promoter.

14. The genetically modified or viral infected plant or cultured plant cells of claim 12, wherein said fusion protein is compartmentalized within a cellular compartment selected from the group consisting of

cytoplasm, endoplasmic reticulum, golgi apparatus, oil bodies, starch bodies, chloroplastids, chloroplasts, chromoplastids, chromoplasts, vacuole, lysosomes, mitochondria, and nucleus.

15. A genetically modified or viral infected plant or cultured plant cells expressing a fusion protein including a recombinant protein and a cellulose binding peptide separated therebetween via a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence.

16. The genetically modified or viral infected plant or cultured plant cells of claim 15, wherein said fusion protein is compartmentalized within cells of said plant or cultured plant cells, so as to be sequestered from cell walls of said cells of said plant or cultured plant cells.

17. The genetically modified or viral infected plant or cultured plant cells of claim 15, wherein said fusion protein is compartmentalized within a cellular compartment selected from the group consisting of cytoplasm, endoplasmic reticulum, golgi apparatus, oil bodies, starch bodies, chloroplastids, chloroplasts, chromoplastids, chromoplasts, vacuole, lysosomes, mitochondria, and nucleus.

18. The genetically modified or viral infected plant or cultured plant cells of claim 15, wherein expression of said fusion protein is under a control of a constitutive or tissue specific plant promoter.

19. A composition of matter comprising:

(a) a plant derived cellulosic matter of a plant; and

- (b) a fusion protein including a recombinant protein and a cellulose binding peptide separated therebetween via a unique amino acid sequence recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence, said fusion protein being expressed in said plant and complexed to said plant derived cellulosic matter of said plant by affinity binding via said cellulose binding peptide.

20. A nucleic acid molecule comprising:

- (a) a promoter sequence for directing protein expression in plant cells;
- (b) a heterologous nucleic acid sequence including:
 - (i) a first sequence encoding a cellulose binding peptide;
 - (ii) a second sequence encoding a recombinant protein, wherein said first and second sequences are joined together in frame; and
 - (iii) a third sequence encoding a unique amino acid sequence being recognizable and digestible by a protease or under predetermined cis or trans conditions for digesting a controllable intervening protein sequence, said third sequence being between and in frame with said first and second sequences;

wherein, said heterologous nucleic acid sequence being down stream said promoter sequence, such that expression of said heterologous nucleic acid sequence is effectable by said promoter sequence.

21. The nucleic acid molecule of claim 20, further comprising a sequence element selected from the group consisting of an origin of

replication for propagation in bacterial cells, at least one sequence element for integration into a plant's genome, a polyadenylation recognition sequence, a transcription termination signal, a sequence encoding a translation start site, a sequence encoding a translation stop site, plant RNA virus derived sequences, plant DNA virus derived sequences, tumor inducing (Ti) plasmid derived sequences, and a transposable element derived sequence.

22. A nucleic acid molecule comprising:

- (a) a promoter sequence for directing protein expression in plant cells;
- (b) a heterologous nucleic acid sequence including:
 - (i) a first sequence encoding a cellulose binding peptide;
 - (ii) a second sequence encoding a recombinant protein, wherein said first and second sequences are joined together in frame; and
 - (iii) a third sequence encoding a signal peptide for directing a protein to a cellular compartment, said third sequence being upstream and in frame with said first and second sequences;

wherein, said heterologous nucleic acid sequence being down stream said promoter sequence, such that expression of said heterologous nucleic acid sequence is effectable by said promoter sequence.

23. The nucleic acid molecule of claim 22, wherein said heterologous nucleic acid sequence further includes

- (iv) a fourth sequence encoding a unique amino acid sequence being recognizable and digestible by a protease or under predetermined cis or trans conditions

for digesting a controllable intervening protein sequence, said fourth sequence being between and in frame with said first and second sequences.

24. The nucleic acid molecule of claim 22, further comprising a sequence element selected from the group consisting of an origin of replication for propagation in bacterial cells, at least one sequence element for integration into a plant's genome, a polyadenylation recognition sequence, a transcription termination signal, a sequence encoding a translation start site, a sequence encoding a translation stop site, plant RNA virus derived sequences, plant DNA virus derived sequences, tumor inducing (Ti) plasmid derived sequences, and a transposable element derived sequence.

25. The nucleic acid molecule of claim 22, wherein said cellular compartment is selected from the group consisting of cytoplasm, endoplasmic reticulum, golgi apparatus, oil bodies, starch bodies, chloroplastids, chloroplasts, chromoplastids, chromoplasts, vacuole, lysosomes, mitochondria, and nucleus.

ABSTRACT OF THE DISCLOSURE

A process of expressing a recombinant protein in a plant and of isolating the recombinant protein from the plant, the process is effected by (a) providing a plant, a plant derived tissue or cultured plant cells expressing a fusion protein including the recombinant protein and a cellulose binding peptide being fused thereto, the fusion protein being compartmentalized within cells of the plant, plant derived tissue or cultured plant cells, so as to be sequestered from cell walls of the cells of the plant, plant derived tissue or cultured plant cells; (b) homogenizing the plant, plant derived tissue or cultured plant cells, so as to bring into contact the fusion protein with a cellulosic matter of the plant, plant derived tissue or cultured plant cells, to thereby effect affinity binding of the fusion protein via the cellulose binding peptide to the cellulosic matter, thereby obtaining a fusion protein cellulosic matter complex; and (c) isolating the fusion protein cellulosic matter complex.

Combined Declaration For Patent Application and Power of Attorney

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled PROCESS OF EXPRESSING AND ISOLATING RECOMBINANT PROTEINS AND RECOMBINANT PROTEIN PRODUCTS FROM PLANTS, PLANT DERIVED TISSUES OR CULTURED PLANT CELLS, the specification of which

(check one) ☒ is attached hereto

☐ was filed on _____ as Application Serial No. _____ and was amended on _____. I hereby state that I

have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a)

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

NONE

(number) (Country) (Day, Month, Year Filed)

☐ ☐

Yes No

(number) (Country) (Day, Month, Year Filed)

☐ ☐

Yes No

(number) (Country) (Day, Month, Year Filed)

☐ ☐

Yes No

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States Application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)

(Filing Date)

Status

(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

Status

(patented, pending, abandoned)

I hereby appoint the following attorneys, with full power of substitution, association, and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

Mark M. Friedman Registration No. 33,883

Address all Correspondence to:

DR. MARK FRIEDMAN LTD.
c/o ANTHONY CASTORINA
2001 JEFFERSON DAVIS HIGHWAY
SUITE 207
ARLINGTON, VIRGINIA 22202

Direct all telephone calls & faxes to:
ROBERT SHEINBEIN
Phone (703) 415-1581
Fax (703) 415-4864

Continuation of Combined Declaration For Patent Application and Power of Attorney

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statement may jeopardize the validity of the application of any patent issued thereon.

*FULL NAME OF SOLE OR FIRST INVENTOR ZIV SHANI	INVENTOR'S SIGNATURE <i>Ziv Shani</i>	DATE 25/5/99
RESIDENCE 4 PALDI, 76248 REHOVOT, ISRAEL	CITIZENSHIP ISRAELI	
POST OFFICE ADDRESS 4 PALDI, 76248 REHOVOT, ISRAEL		

*FULL NAME OF SECOND INVENTOR ODED SHOSEYOV	INVENTOR'S SIGNATURE <i>Oded Shoseyov</i>	DATE 25/5/99
RESIDENCE 5 HAEREZ, 72910 KARME YOSEF, ISRAEL	CITIZENSHIP ISRAELI	
POST OFFICE ADDRESS 5 HAEREZ, 72910 KARME YOSEF, ISRAEL		

*FULL NAME OF FOURTH INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE	CITIZENSHIP ISRAELI	
POST OFFICE ADDRESS		

*FULL NAME OF FIFTH INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE	CITIZENSHIP ISRAELI	
POST OFFICE ADDRESS		

*FULL NAME OF SIXTH INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE	CITIZENSHIP ISRAELI	
POST OFFICE ADDRESS		

*FULL NAME OF SEVENTH INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE	CITIZENSHIP ISRAELI	
POST OFFICE ADDRESS		

SMALL BUSINESS CONCERN - NEW APPLICATION

Attorney Docket No. 1079/5

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In RE Application of ZIV SHANI ET AL.

Filed Concurrently Herewith

For PROCESS OF EXPRESSING AND ISOLATING RECOMBINANT PROTEINS AND RECOMBINANT PROTEIN PRODUCTS FROM PLANTS, PLANT DERIVED TISSUES OR CULTURED PLANT CELLSVERIFIED STATEMENT UNDER 37 CFR 1.27
CLAIMING STATUS AS A SMALL ENTITY

To The Commissioner of Patents and Trademarks:

I hereby declare that:

I am the owner of, or an official empowered to act on behalf of, the small business concern identified below:

Name of Concern: CBD TECHNOLOGIES LTD.Address : 2 PEKERIS, 4th FLOOR, PARK TAMAR, P. O. BOX 199, 76100 REHOVOT, ISRAEL

The small business concern identified above, together with its affiliates, employs fewer than 500 persons and qualifies as a small business concern as defined in 37 CFR 1.9(d) for purposes of paying reduced fees under 35 USC § 41(a) and § 41(b) to the Patent and Trademark Office with regard to the above-entitled invention described in the specification filed herewith.

Rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the above entitled invention.

If the rights held by the small business concern are not exclusive, each other party having rights to the invention is listed below, and no rights to the invention are held by any party who could not qualify as a small entity under 37 CFR 1.9(f), namely any person who could not be classified as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

Full Name (Party 1) : NONE

Address : _____

Status : ☐ Individual ☐ Small Business Concern ☐ Nonprofit Organization

Full Name (Party 2) : _____

Address : _____

Status : ☐ Individual ☐ Small Business Concern ☐ Nonprofit Organization

acknowledge the duty under 37 CFR 1.28(b) to file, in this application, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the issue fee due after the date on which status as a small entity is no longer appropriate.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application and any patent issuing thereon.

DR. STANLEY HIRSCH.

Name of Person Signing

Signature

June 7, 1999
DateCapacity of Person Signing: PresidentAddress of Person Signing : 2 PEKERIS, REHOVOT, ISRAEL

972 3 562554

SMALL BUSINESS CONCERN - NEW APPLICATION

Attorney Docket No.: 1079/5

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In RE Application of: ZIV SHANI ET AL.

Filed Concurrently Herewith

For: PROCESS OF EXPRESSING AND ISOLATING RECOMBINANT PROTEINS AND RECOMBINANT PROTEIN PRODUCTS FROM PLANTS, PLANT DERIVED TISSUES OR CULTURED PLANT CELLSVERIFIED STATEMENT UNDER 37 CFR 1.27
CLAIMING STATUS AS A SMALL ENTITY

To The Commissioner of Patents and Trademarks:

I hereby declare that:

I am the owner of, or an official empowered to act on behalf of, the small business concern identified below.

Name of Concern: YISSUM RESEARCH AND DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEMAddress : 46 JABOTINSKI, 91042 JERUSALEM, ISRAEL

The small business concern identified above, together with its affiliates, employs fewer than 500 persons and qualifies as a small business concern as defined in 37 CFR 1.8(d) for purposes of paying reduced fees under 35 USC § 41(a) and § 41(b) to the Patent and Trademark Office with regard to the above-entitled invention described in the specification filed herewith.

Rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the above entitled invention.

If the rights held by the small business concern are not exclusive, each other party having rights to the invention is listed below, and no rights to the invention are held by any party who could not qualify as a small entity under 37 CFR 1.8(f), namely any person who could not be classified as an independent inventor under 37 CFR 1.8(c) if that person had made the invention, or any concern which would not qualify as a small business concern under 37 CFR 1.8(d) or a nonprofit organization under 37 CFR 1.8(e).

Full Name (Party 1) : NONE

Address : _____

Status : ☐ Individual☐ Small Business
Concern☐ Nonprofit
Organization

Full Name (Party 2) : _____

Address : _____

Status : ☐ Individual☐ Small Business
Concern☐ Nonprofit
Organization

I acknowledge the duty under 37 CFR 1.28(b) to file, in this application, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the issue fee due after the date on which status as a small entity is no longer appropriate.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application and any patent issuing thereon.

MORDEHAI PERCHUTER

Name of Person Signing

Signature

JUNE 3, 1999

Date

Capacity of Person Signing: MANAGING DIRECTOR & CEOAddress of Person Signing: 46 JABOTINSKI ST JERUSALEM 91042

Yissum Research Development Co.
of the Hebrew University of Jerusalem
P.O.B. 4279, Jerusalem, 91042, ISRAEL

FIGURE 1

